# Neural precursor and stem cells

Patent number:

EP1529838

**Publication date:** 

2005-05-11

Inventor:

BOSIO ANDREAS DR (DE); CREMER HAROLD (FR);

PENNARTZ SANDRA (DE)

Applicant:

MEMOREC BIOTEC GMBH (DE)

Classification:

- international:

C12N5/06; G01N33/53

- european:

C12N5/06B8P; G01N33/68V2

Application number: EP20030025506 20031106 Priority number(s): EP20030025506 20031106 Cited documents:

EP1354943 XP00227572

> XP00222350 XP00103392

XP00902890

Report a data error he

## Abstract of EP1529838

A cell population comprising at least 5% neural stem cells, the stem cells being characterized by an expression of ASCT2 or KIAA0152, is new. - Independent claims are also included for the following: - (1) a method for isolating the cell population cited above; - (2) a medicament comprising the above cell population; and - (3) a monoclonal antiboc directed against ASCT2. - ACTIVITY - Neuroprotective; Nootropic; Antiparkinsonian; Cerebroprotective; Vasotropic; No biological data given - MECHANISM OF ACTION - Cell Therapy.

Data supplied from the esp@cenet database - Worldwide



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11) EP 1 529 838 A1

(12)

# **EUROPÄISCHE PATENTANMELDUNG**

(43) Veröffentlichungstag: 11.05.2005 Patentblatt 2005/19 (51) Int Cl.7: C12N 5/06, G01N 33/53

(21) Anmeldenummer: 03025506.1

(22) Anmeldetag: 06.11.2003

(84) Benannte Vertragsstaaten:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IT LI LU MC NL PT RO SE SI SK TR
Benannte Erstreckungsstaaten:

AL LT LV MK

(71) Anmelder: Memorec Biotec GmbH 50829 Köln (DE)

(72) Erfinder:

 Bosio, Andreas, Dr. 50931 Köln (DE)  Cremer, Harold 13400 Marseille (FR)

 Pennartz, Sandra 50670 Köln (DE)

(74) Vertreter: Schreiber, Christoph, Dr. et al Patentanwälte von Kreisler Selting Werner, Postfach 10 22 41 50462 Köln (DE)

(54) Neurale Vorläufer- und Stammzellen

(57) Zellpopulation, dadurch gekennzeichnet, dass mindestens 5% der Zellen neurale Vorläuferzellen sind, die wenigstens einen der in **Liste A** oder **Liste B** aufgeführten Marker aufweisen.

25

2

#### Beschreibung

[0001] Die vorliegende Erfindung betrifft Zellpopulationen von neuralen Vorläuferzellen bzw. neuralen Stammzellen sowie Verfahren zur Isolierung entsprechender Zellen.

[0002] Der Ausgangspunkt für die Generierung der über tausend verschiedenen neuronalen und glialen Zelltypen des Nervensystems von Vertebraten sind multipotente, neurale Stammzellen des embryonalen Neuroepitheliums (Williams, B. P., Read, J. & Price, J. (1991): The generation of neurons and oligodendrocytes from a common precursor cell. Neuron 7(4), 685-93), (Davis, A. A. & Temple, S. (1994): A self-renewing multipotential stem cell in embryonic rat cerebral cortex. Nature 372(6503), 263-6), (Weiss, S., Dunne, C., Hewson, J., Wohl, C., Wheatley, M., Peterson, A. C. & Reynolds, B. A. (1996): Multipotent CNS stem cells are present in the adult mammalian spinal cord and ventricular neuroaxis. J Neurosci 16(23), 7599-609).

[0003] In den vergangenen Jahren wurde durch verschiedene Arbeitsgruppen gezeigt, dass solche sich selbst erneuernden, multipotenten Vorläuferzellen nicht nur während der Entwicklung, sondern auch im adulten Gehirn zu finden sind (Gage, F. H. (2000): Mammalian neural stem cells. *Science* 287(5457), 1433-8). Vor allem um die lateralen Ventrikel des Vorderhirns findet die Bildung von neuralen Vorläuferzellen lebenslang statt. Diese wandern hauptsächlich, wenn auch nicht exklusiv, in den Bulbus olfaktorius, um dort in GABA-erge Interneurone zu differenzieren.

[0004] Über die genaue Lokalisation der multipotenten Stammzellen, die dieser sekundären Neurogenese zugrunde liegen, wird derzeit noch spekuliert: Johansson et al. beschrieben ependymale Zellen entlang des Lumen der adulten, ventrikulären Zone mit den Eigenschaften multipotenter Stammzellen (Johansson, C. B., Svensson, M., Wallstedt, L., Janson, A. M. & Frisen, J. (1999b): Neural stem cells in the adult human brain. Exp. Cell Res 253(2), 733-6), während Doetsch et al. Astrocyten der subventrikulären Zone als multipotente Stammzellen identifizierten (Doetsch, F., Caille, I., Lim, D. A., Garcia-Verdugo, J. M. & Alvarez-Buylla, A. (1999): Subventricular zone astrocytes are neural stem cells in the adult mammalian brain. Cell 97(6), 703-16). Eine absolut eindeutige Identifizierung dieser adulten Stammzellen in vivo ist jedoch bis heute, hauptsächlich mangels geeigneter Marker, nicht gelungen.

[0005] Neben ihrer Bedeutung im olfaktorischen System ist das therapeutische Potential der adulten Stammzellen von besonderem Interesse. Aufgrund ihrer Multipotenz weisen neurale Stammzellen bemerkenswerte Formbarkeit auf und könnten daher durch Zusatz von verschiedenen Faktoren zur Erzeugung verschiedener Neuronentypen eingesetzt werden. Die anschließende Transplantation der so entwickelten spezialisierten Zellen könnte zur Behandlung von neurologischen Krankheiten Alzheimer, Parkinson, Folgen von

Schädel-Hirn-Traumata und Schlaganfall beitragen. Voraussetzung dafür ist die Charakterisierung der verschiedenen, neuralen Differenzierungsstufen sowie die Identifizierung der Faktoren, die die Differenzierungsprogramme der Stammzellen steuern. Gegenüber den embryonalen Stammzellen haben die adulten den Vorteil, dass sie erstens keine abstoßende Immunreaktion auslösen würden, weil sie dem Körper des Patienten entstammen, folglich ihre Transplantation ohne Immunsuppression erfolgen könnte, und zweitens ihre Gewinnung ethisch unbedenklich ist.

[0006] Die Erforschung der Eigenschaften neuraler Stammzellen und embronaler Stammzellen des Menschen ist aus ethischen Aspekten praktisch nicht oder nur sehr eingeschränkt möglich. Daher wurden alle explorativen Arbeiten ausgehend von Mäusen und Mauszellen durchgeführt. Wie bereits beschrieben war die Isolierung von neuralen Stammzellen bisher nicht möglich, da dieser Zelltyp nicht eindeutig charakterisiert war und keine geeigneten Marker zur Identifizierung und Anreicherung zur Verfügung standen.

[0007] Aufgabe der vorliegenden Erfindung war es daher Verfahren zu entwickeln, die eine Isolation von neuralen Vorläuferzellen und neuralen Stammzellen erlauben und entsprechende Zellpopulation, enthaltend diese Zelle bereitzustellen.

[0008] Erfindungsgemäß wird die Aufgabe gelöst durch die Identifizierung von Markern, die entsprechende Zellen aufweisen.

[0009] Marker ist ein Gen, das mit Hilfe der Serial Analysis of Genexpression (SAGE) in entsprechenden Zellen gefunden wird.

[0010] Methodisch beruht SAGE auf der Isolierung von 14 bp großen DNA Fragmenten (Tags), die jeweils charakteristisch für eine mRNA-Spezies sind. Die Tags, repräsentativ für alle in der zu untersuchenden Zelle vorliegenden mRNA Moleküle, werden zu langen Polymeren verbunden, die im letzten Schritt der Methode sequenziert werden. Die Frequenz, mit der ein Tag sequenziert wird, ist direkt proportional zur Kopienzahl der mRNA-Moleküle im untersuchten Ausgangsmaterial (Velculescu, V. E., Zhang, L., Vogelstein, B. & Kinzler, K. W. (1995): Serial analysis of gene expression. Science 270(5235), 484-7). Durch die computerunterstützte Auswertung der Sequenzdaten entsteht ein digitales Expressionsprofil, das beliebig oft und ohne zusätzliche Laborarbeit mit Expressionsprofilen anderer Gewebe verglichen werden kann (Meta-Analyse).

[0011] Den so identifizierten Gene sind eindeutigen Nummern zugeordnet, die beispielsweise als SAGEmap von National Center for Biotechnology Information (NCBI) bereitgestellt werden (www.ncbi.nlm.nin.gov/SAGE).

[0012] Gegenstand der Erfindung sind zum einen Zellpopulationen, bei denen mindestens 5% der Zellen neurale Vorläuferzellen sind, die wenigstens einen der in Liste A oder Liste B aufgeführten Marker aufweisen. [0013] Bevorzugt weisen entsprechende neurale Vor-

läuferzellen wenigstens zwei, drei, vier oder fünf der in Liste A oder B aufgeführten Marker auf.

3

[0014] In bevorzugten Ausführungsformen weisen entsprechende neurale Vorläuferzellen keinen der in Liste C aufgeführten Marker auf.

[0015] Bevorzugt ist der Gehalt an neuralen Vorläuferzellen in der Zellpopulation hoch, d.h. mindestens 10%, bevorzugt mindestens 25%, noch mehr bevorzugt mehr als 50% und am meisten bevorzugt über 90%.

[0016] Entsprechende neurale Vorläuferzellen sind vorzugsweise aus Hirngewebe erhältlich.

[0017] In einer Ausführungsform handelt es sich dabei um eine murine Zellpopulation.

[0018] Gegenstand der Erfindung ist auch ein Verfahren zur Isolierung einer entsprechenden Zellpopulation mit folgenden Schritten:

## entweder

- Entnahme einer Probe aus dem Hirn
- Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker

#### oder

- Differenzierung von embryonalen Stammzellen zu neuralen Vorläuferzellen,
- Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker

## oder

- Trans-Differenzierung von adulten, nicht neuralen Stammzellen zu neuralen Vorläuferzellen,
- Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker

## oder

- Differenzierung von adulten, neuralen Stammzellen zu neuralen Vorläuferzellen,
- Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker

## oder

- Differenzierung von immortalisierten Zellen zu neuralen Vorläuferzellen,
- Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker.

[0019] "Unter Verwendung der angegebenen Marker" bedeutet, dass die Zellen isoliert werden, die positiv für mindestens einen der Marker aus der Liste A und B sind, wobei mehrere positive Marker und die Abwesenheit von Markern der Liste C bevorzugt werden. Die Isolierung kann beispielsweise durch FACS Analyse erfolierung kann beispielsweise durch FACS Analyse er

gen. Die durch die Verfahren erhältlichen Zellen sind ebenfalls Gegenstand der Erfindung.

4

[0020] Ein weiterer Gegenstand der Erfindung ist die Verwendung mindestens eines Markers ausgewählt aus der Liste A oder Liste B zu Identifizierung oder Isolierung von neuralen Vorläuferzellen.

[0021] Gegenstand ist weiterhin ein Antikörper gegen einen Marker aus der Liste A, B oder C, ein Diagnostikmittel enthaltend mindestens einen, bevorzugt zwei oder mehr Substanzen zur Erkennung der Marker der Liste A, B oder C sowie ein Arzneimittel enthaltend die erfindungsgemäße Zellpopulation.

[0022] Solche Arzneimittel könnten wie oben dargestellt zur Behandlung von neurologischen Krankheiten wie Alzheimer, Parkinson, Folgen von Schädelhirntraumata oder Schlaganfall eingesetzt werden.

[0023] Ein weiterer Gegenstand ist eine Zellpopulation, bei der mindestens 5% der Zellen neurale Stammzellen sind, die wenigstens einen der in Liste D oder Liste E aufgeführten Marker aufweisen.

[0024] Vorzugsweise weisen entsprechende neurale Stammzellen mindestens zwei, bevorzugt mindestens drei, mindestens vier und noch mehr bevorzugt mindestens fünf der in Liste D oder Liste E aufgeführten Marker auf.

[0025] In besonders bevorzugten Ausführungsformen weisen entsprechende neurale Stammzellen keinen der in Liste A oder Liste C aufgeführten Marker auf. [0026] Der Gehalt an neuralen Stammzellen in der Zellpopulation ist möglichst hoch, bevorzugt mindestes 10%, mehr bevorzugt mindestes 25%, mindestens 50%, und am meisten bevorzugt mindestens 90%.

[0027] Entsprechende Zellpopulation sind aus Hirngewebe erhältlich. In einer Ausführungsform handelt es sich um eine murine Zellpopulation.

[0028] Gegenstand ist weiterhin ein Verfahren zur Isolierung der Zellpopulation. Dies ist erhältlich entweder durch

- 40 Entnahme einer Probe aus dem Hirn
  - Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker

# 45 oder

- Differenzierung von embryonalen Stammzellen zu neuralen Stammzellen,
- Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker

## oder

- 55 Trans-Differenzierung von adulten, nicht neuralen Stammzellen zu neuralen Stammzellen,
  - Isolieren der neuralen Stammzellen unter Verwen-

ten gezeigt.

5

dung der angegebenen Marker

#### oder

- De-Differenzierung von adulten, neuralen Vorläuferzellen zu neuralen Stammzellen.
- Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker

#### oder

- Differenzierung von immortalisierten Zellen zu neuralen Stammzellen,
- Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker.

[0029] Die Isolierung erfolgt wie oben bei den neuralen Vorläuferzellen angegeben. Auch die auf diesem Wege erhältlichen neuralen Stammzellen sind Gegenstand der Erfindung.

[0030] Gegenstand der Erfindung ist weiterhin ein Antikörper gegen einen Marker aus der Liste D, E, ein Diagnostikmittel enthaltend mindestens einen, bevorzugt zwei oder mehr Substanzen zur Erkennung der Marker der Liste D, E, A oder C sowie ein Arzneimittel enthaltend die erfindungsgemäße Zellpopulation.

[0031] Solche Arzneimitteln können wie dargestellt zur Behandlung von neuronalen Krankheiten wie Alzheimer, Parkinson, Folgen von Schädelhirntraumata oder Schlaganfall eingesetzt werden.

## Beispiele

# A. Isolierung von embryonaler Stammzellen

[0032] Murine embryonale Stammzellen proliferieren klonal in vitro und sind aus diesem Grunde in großer Menge und hochreiner Form isolierbar. Nach dem Stand der Technik werden diese in Anwesenheit von LIF auf primären embryonalen Fibroblasten gehalten und regelmäßig durch die Generierung von hochgradig keimbahnkompetenten chimären Mäusen auf ihre Qualität überprüft. Unter normalen Kulturbedingungen beträgt das Verhältnis ES-Zellen zu kontaminierenden Fibroblasten etwa 200:1. Um auch diese minoritäre Komponente zu eliminieren, wurden die ES-Zellen vor der RNA-Päparation für zwei Passagen (vier Tage) auf gelatinisierten Kulturplatten bei erhöhter LIF-Konzentration gehalten. Dies ermöglicht eine Reduktion der kontaminierenden Fibroblasten auf etwa 0,01% der Gesamtpopulation.

B. Isolierung von neuronalen Vorläuferzellen aus dem adulten Mausgehirn.

[0033] In der subventrikulären Zone des adulten Vor-

derhirns von Vertebraten werden permanent große Mengen von neuralen Vorläuferzellen gebildet (wahrscheinlich < 50000 Zellen/ Tag). Diese Zellen benutzen einen präzise definierten Migrationsweg und eine spezielle Form der Translokation (*Chain migration*) um in den Bulbus olfaktorius zu gelangen. Im Bulbus olfaktorius angelangt differenzieren diese Vorläuferzellen normalerweise in inhibitorische (GABA-erge) Interneurone. Unter bestimmten experimentellen Bedingungen wurde ihre Differenzierung in Oligodendrozyten und Astrozy-

6

[0034] Neurale Vorläufer, die einen Differenzierungszustand zwischen einer neuralen Stammzelle und einem terminal differenzierten Neuron repräsentieren, exprimieren spezifisch eine Form des neuralen Zelladhäsionsmoleküls NCAM, die eine spezielle post-translationelle Modifikation aufweist. Diese Modifikation besteht aus der Glykosylierung des Proteins mit a-2,8 verknüpfter Polysialylsäure (PSA). Ein spezifischer Antikörper gegen dieses Glykoepitop (Chazal et al., 2000) erlaubte die hochreine Isolierung der Zielpopulation aus dissozierten Vorderhirngewebe durch FACS (Fluorescence Activated Cell Sorting).

## <sup>25</sup> C. Molekulargenetische Analyse

[0035] Embryonale Stammzellen und neuronale Vorläuferzellen wurden in einem genomweiten Screen mit der Methode SAGE (Serial Analysis of Gene Expression) analysiert.

[0036] Die Genexpressionsprofile der beiden Zell-Populationen wurden unter Anwendung bioinformatischer Verfahrensweisen mit Maus-Hirn-SAGE-Datenbanken verglichen, um molekulare Marker zu identifizieren, die charakteristisch für embryonale Stammzellen und neuronale Vorläuferzellen sind.

[0037] Mit Hilfe der Microarray technologie wurde die Expression der Gene bestätigt.

[0038] Durch in situ-Hybridisierung in Maushirn und an embryonalen Stammzellen wurde die zelluläre Lokalisation einiger der identifizierten Gene bestimmt. Diese Ergebnisse belegen, dass spezifische Markergene identifiziert werden konnten.

45 Liste A: Positivmarker neurale Vorläuferzellen (1.) und Negativmarker 2 neurale Stammzellen;

ES-Zellen -; PSA-NCAM +; Adult brain -

# 50 [0039]

	Mm.8884	nuclear factor of kappa light chain gene				
		enhancer in B-cells inhibitor, alpha				
	Mm.8180	lymphocyte antigen 6 complex, locus A				
55	Mm.6238	SRY-box containing gene 11				
	Mm.517	(Manual) Manic fringe protein, putative				
		secreted glycosyltransferase, notch				
		modulator				

7

Mm.4919 Mm.4727	DNA segment, human D4S114 seizure related gene 6		Liste B: Pos	itivmarker neurale Vorläuferzellen (2.);
Mm.45769	ESTs		ES-Zellen -/-	-; PSA-NCAM +; Adult brain -
Mm.44490	RIKEN cDNA 6330415M09 gene			
Mm.42948	peroxiredoxin 2	5	[0040]	
Mm.4022	RIKEN cDNA 1110033C18 gene			
Mm.3940	lethal giant larvae homolog		Mm.911	high mobility group nucleosomal bin-
Mm.37835	ribosomal protein L7			ding domain 2
Mm.3779	RIKEN cDNA 2300006C11 gene		Mm.89136	H3 histone, family 3A
Mm.340	high mobility group box 3	10	Mm.741	fatty acid binding protein 5, epidermal
Mm.32902	ESTs, Weakly similar to S26689 hypo-		Mm.7286	C-terminal binding protein 1
NA 0000	thetical protein hc1 - mouse		Mm.7141	proliferating cell nuclear antigen
Mm.3268	ubiquitin-conjugating enzyme E2I		Mm.6840	RIKEN cDNA 5730507C05 gene
Mm.31436	myeloid ecotropic viral integration site-	15	Mm.6787	splicing factor, arginine/serine-rich 3
Mm.297	related gene 1 actin, beta, cytoplasmic	,5	Mm.6417	(SRp20) CD24a antigen
Mm.29558	expressed sequence Al426163		Mm.6343	nucleophosmin 1
Mm.29014	T-cell lymphoma invasion and metasta-		Mm.482	Jun oncogene
141111.20014	sis 2		Mm.43871	expressed sequence AW046487
Mm.28842	chloride channel 3	20	Mm.43213	RIKEN cDNA 9030402K04 gene
Mm.28824	Mus musculus, clone IMAGE:4504748,		Mm.42767	ribosomal protein S17
	mRNA		Mm.4269	transcription factor 4
Mm.28275	RNA binding motif protein, X chromoso-		Mm.40715	RIKEN cDNA 1110038H03 gene
	me		Mm.40715	RIKEN cDNA 1110038H03 gene
Mm.28149	RIKEN cDNA 3110003A17 gene	25	Mm.4071	laminin receptor 1 (67kD, ribosomal pro-
Mm.28148	chromobox homolog 3 (Drosophila HP1			tein SA)
	gamma)		Mm.4025	nuclear factor I/B
Mm.27816	hexosaminidase B		Mm.372	ribosomal protein S26
Mm.2769	MARCKS-like protein		Mm.3487	ribosomal protein L30
Mm.22171	calponin 3, acidic	30	Mm.3381	ribosomal protein S8
Mm.220923	RIKEN cDNA 6530406007 gene		Mm.31051	RIKEN cDNA 2610003J05 gene
Mm.21740	heterogeneous nuclear ribonucleopro-		Mm.30120	ribosomal protein S27-like
	tein H1		Mm.30011	ribosomal protein S23
Mm.206085	expressed sequence Al854782	05	Mm.29911	RIKEN cDNA 3200001M24 gene
Mm.205996	EST AA087124	35	Mm.2966	isocitrate dehydrogenase 2 (NADP+),
Mm.200858 Mm.199500	RIKEN cDNA 2410129E14 gene		Mm.29580	mitochondrial
Mm.195901	expressed sequence Al844617 ribosomal protein L35a		WITT.29560	superiorcervical ganglia, neural specific 10
Mm. 194965	EST		Mm.2958	expressed sequence Al843786
Mm.19101	DEAD (aspartate-glutamate-alanine-	40	Mm.28985	ribosomal protein L27
141111.13101	aspartate) box polypeptide 5	,,	Mm.28869	ESTs
Mm.19016	drebrin 1		Mm.27927	heterogeneous nuclear ribonucleopro-
Mm. 18789	SRY-box containing gene 4			tein A1
Mm.186740	ESTs		Mm.27669	small nuclear ribonucleoprotein E
Mm.18516	H3 histone, family 3B	45	Mm.2756.	high mobility group nucleosomal bin-
Mm.181959	early growth response 1			ding domain 1
Mm.181847	prefoldin 5		Mm.27141	Rac GTPase-activating protein 1
Mm.16421	high mobility group box 1		Mm.2591	RNA binding motif protein 3
Mm.15534	interleukin 1 alpha		Mm.24083	Mus musculus, Similar to TAR DNA bin-
Mm.13725	Paneth cell enhanced expression	50		ding protein, clone MGC: 19284
Mm.12871	doublecortin			IMAGE:4016437, mRNA, complete cds
Mm.127662	ESTs		Mm.219668	RIKEN cDNA 2610209F03 gene
Mm.12412	Mus musculus, Similar to RIKEN cDNA 2810407E23 gene, clone IMAGE:		Mm.21841	splicing factor, arginine/serine-rich 2 (SC-35)
	4489006, mRNA, partial cds	55	Mm.218240	Mus musculus, clone IMAGE:5342828, mRNA, partial cds
			Mm.21740	heterogeneous nuclear ribonucleoprotein H1

10

Mm.213020 (Manual) 60S ribosomal protein L32 Mm.6660 small inducible cytokine A27 (RPL32) Mm.6586 Mus musculus, clone MGC:6299 IMAGE: Mm.2115 heterogeneous nuclear ribonucleopro-2654341, mRNA, complete cds tein U Mm.6565 FK506 binding protein 8 (38 kDa) Mm.196611 synapsin I Mm.65337 Mus musculus, clone MGC:28924 Mm.19187 prothymosin alpha IMAGE:3481738, mRNA, complete cds Mm.18789 SRY-box containing gene 4 Mm.648 prion protein Mm.186499 ESTs, Weakly similar to immunoglobulin Mm.638 **ESTs** superfamily containing leucinerich re-Mm.544 phosphoprotein enriched in astrocytes 15 peat Mm.5264 ESTs, Highly similar to FEZ1\_RAT FA-Mm.18516 SCICULATION AND ELONGATION H3 histone, family 3B Mm.180873 RIKEN cDNA 2510019J09 gene PROTEIN ZETA 1 (ZYGIN I) Mm.1775 hematological and neurological expres-Mm.5259 (Manual assignment) probably myelinsed sequence 1 associated oligodendrocyte basic protein Mm.1703 tubulin, beta 5 15 MOBP Mm.16775 copine 6 ribosomal protein S24 Mm.5249 Mm.16767 heterogeneous nuclear ribonucleopro-Mm.52 RIKEN cDNA 1810033A19 gene tein A2/B1 Mm.5195 complexin 1 Mm. 16596 B-cell translocation gene 1, anti-prolife-Mm.5153 neurotensin receptor 2 Mm.5023 Purkinje cell protein 4 Mm.148973 RIKEN cDNA 3010025E17 gene Mm.4923 **ESTs** Mm.142872 heterogeneous nuclear ribonucleopro-Mm.4921 glutamate receptor, ionotropic, AMPA2 tein K (alpha 2) Mm.142729 thymosin, beta 4, X chromosome Mm.4920 glutamate receptor, ionotropic, AMPA1 Mm.140380 ribosomal protein L23 25 (alpha 1) Mm.140 protein phosphatase 1, regulatory (inhi-Mm.4870 synaptosomal-associated protein, 91 bitor) subunit 14B kDa Mm. 12858 eukaryotic translation initiation factor Mm.4857 calcium/calmodulin-dependent protein kinase II, beta kinesin heavy chain member 1A 30 Mm.4762 Liste C: Negativmarker 1 neurale Stammzellen und Mm.4705 (Manual) probably in far 3'-UTR of complexin-2 cDNA Negativmarker neurale Vorläuferzellen; Mm.46764 RIKEN cDNA 4833409J18 gene ES-Zellen -; PSA-NCAM -; Adult brain + Mm.4657 amyloid beta (A4) precursor protein-bin-35 ding, family A, member 2 [0041] Mm.4651 kinesin-associated protein 3 RIKEN cDNA 1200016B17 gene Mm.45951 Mm.98 ATPase, Na+/K+ transporting, beta 1 poproteasome (prosome, macropain) subu-Mm.4550 nit, beta type 6 lypeptide Mm.9745 lactate dehydrogenase 2, B chain Mm.4550 ATPase, Na+/K+ transporting, beta 1 po-Mm.970 creatine kinase, mitochondrial 1, ubiquilypeptide Mm.4537 NADH dehydrogenase (ubiquinone) 1 Mm.891 kinesin family member C2 beta subcomplex, 9 Mm.88833 Mus musculus strain ILS K-CI cotrans-Mm.44355 RIKEN cDNA 6430514L14 gene porter (Slc12a5) mRNA, complete cds 45 Mm.4435 synaptosomal-associated protein, 25 Mm.87027 BM88 antigen kDa Mm.8688 RIKEN cDNA 0610011B04 gene Mm.44244 open reading frame 12 Mm.86654 microtubule-associated protein 6 Mm.44107 **ESTs** Mm.848 testis expressed gene 261 Mm.44101 Mus musculus, ATPase, Na+K+ trans-Mm.806 CD 81 antigen 50 porting, alpha 3 subunit, clone MGC: Mm.80123 ESTs, Weakly similar to simple repeat se-27631 IMAGE:4506376, mRNA, complequence-containing transcript Mm.7729 aldolase 3, C isoform Mm.4383 myc box dependent interacting protein 1 Mm.7420 tubulin, beta 4 Mm.43786 cytochrome c oxidase, subunit VIIc Mm.7363 beta-spectrin 3 Mm.43749 RIKEN cDNA 3100001N19 gene Mm.726 basigin Mm.43721 small nuclear ribonucleoprotein N Mm.7089 necdin Mm.43587 hippocalcin Mm.667 glutathione S-transferase, mu 5 Mm.43415 cytochrome c oxidase, subunit VI a, po-

11

	lypeptide 1		Mm.3974	ubiquitin specific protease 4 (proto-onco-
Mm.4339	laminin, alpha 5			gene)
Mm.43330	RIKEN cDNA 0610025G13 gene		Mm.39548	expressed sequence Al839779
Mm.43278	olfactomedin 1		Mm.3951	thymus cell antigen 1, theta
Mm.43278	olfactomedin 1	5	Mm.3915	myelin-associated oligodendrocytic ba-
Mm.4296	synovial sarcoma translocation, Chromo-			sic protein
	some 18		Mm.39040	myelin and lymphocyte protein, T-cell dif-
Mm.42949	RIKEN cDNA 1110012005 gene			ferentiation protein
Mm.42948	peroxiredoxin 2		Mm.38994	RIKEN cDNA 2600001N01 gene
Mm.42829	selenoprotein W, muscle 1	10	Mm.38993	calsyntenin 1
Mm.4266	integral membrane protein 2B		Mm.38551	calcium binding protein 1
Mm.4266	integral membrane protein 2B		Mm.38469	amyloid beta (A4) precursor protein-bin-
Mm.4263	cystatin C			ding, family B, member 1
Mm.425	histidine triad nucleotide binding protein		Mm.38438	RIKEN cDNA 1200009K17 gene
Mm.42255	ATPase, Ca++ transporting, cardiac	15	Mm.38421	(Manual assignment) ATPase, Na+K+
	muscle, slow twitch 2			transporting, alpha polypeptide
Mm.41926	NADH dehydrogenase (ubiquinone) 1 al-		Mm.38421	(Manual assignment) ATPase, Na+K+
	pha subcomplex, 4			transporting, alpha polypeptide
Mm.41925	RIKEN cDNA 1810034B16 gene		Mm.3840	flotillin 2
Mm.41918	RIKEN cDNA 1110063G11 gene	20	Mm.38248	sialyltransferase 9 (CMP-NeuAc:lacto-
Mm.41911	cytochrome P450, 46 (cholesterol 24-hy-			sylceramide alpha-2,3-sialyltransferase)
	droxylase)		Mm.38036	ESTs, Moderately similar to
Mm.41893	RIKEN cDNA 6330408G06 gene			NX1A_MOUSE_2
Mm.41791	glycoprotein m6b		Mm.38036	ESTs, Moderately similar to
Mm.41752	expressed sequence Al847934	25		NX1A_MOUSE 2
Mm.41735	RIKEN cDNA 2300004C15 gene		Mm.37462	ESTs, Weakly similar to CA11 RAT COL-
Mm.41719	RIKEN cDNA 2610507A21 gene			LAGEN ALPHA 1(I) CHAIN
Mm.41711	Mus musculus, clone IMAGE:3499845,		Mm.37214	transferrin
	mRNA, partial cds		Mm.36275	DNA segment, Chr 11, Brigham & Wo-
Mm.41694	ESTs	30		men's Genetics 0517 expressed
Mm.41692	ESTs, Weakly similar to F59F4.2.p		Mm.3624	guanylate kinase 1
Mm.41642	regulator of G-protein signaling 4		Mm.35837	RIKEN cDNA 2510006D16 gene
Mm.41630	RIKEN cDNA 0710001E10 gene		Mm.35837	RIKEN cDNA 2510006D16 gene
Mm.41604	ESTs, Weakly similar to VAV3_MOUSE		Mm.3544	calcium channel, voltage-dependent, be-
	VAV-3 PROTEIN	35		ta 3 subunit
Mm.41603	expressed sequence Al891706		Mm.35439	secreted acidic cysteine rich glycoprotein
Mm.41603	expressed sequence Al891706		Mm.35270	Ly6/neurotoxin 1
Mm.41602	RIKEN cDNA 3110050007 gene		Mm.3479	ATPase, H+ transporting, lysosomal
Mm.41602	RIKEN cDNA 3110050007 gene			21kDa, V0 subunit B
Mm.4137	chromogranin A	40	Mm.34695	actin related protein 2/3 complex, subunit
Mm.41354	ESTs			1A (41 kDa)
Mm.41277	RIKEN cDNA 1110020M21 gene		Mm.34246	calmodulin 1
Mm.41248	ESTs		Mm.3363	prosaposin
Mm.41190	RIKEN cDNA 1700112L09 gene		Mm.3360	tyrosine 3-monooxygenase/tryptophan
Mm.40863	expressed sequence AW049870	45		5-monooxygenase activation protein, ze-
Mm.40738	RIKEN cDNA 2900072M03 gene			ta polypeptide
Mm.40621	ESTs, Moderately similar to		Mm.33117	ESTs
	Y552_HUMAN HYPOTHETICAL PRO-		Mm.3308	tyrosine 3-monooxygenase/tryptophan
	TEIN KIAA0552			5-monooxygenase activation protein, eta
Mm.40472	expressed sequence AI835002	50		polypeptide
Mm.40443	RIKEN cDNA 4930488B01 gene		Mm.3292	glutamate receptor, ionotropic, NMDA1
Mm.40124	phosphodiesterase 10A			(zeta 1)
Mm.40059	ESTs, Weakly similar to SP62 MOUSE		Mm.3229	ribosomal protein L26
	SPLICEOSOME ASSOCIATED PROTE-		Mm.32191	gamma-aminobutyric acid (GABA-B) re-
	IN 62	55		ceptor, 1
Mm.39857	RIKEN cDNA 2900074L19 gene		Mm.31395	carboxypeptidase E
Mm.39803	expressed sequence Al841080		Mm.3123	cornichon-like (Drosophila)
Mm.39752	RIKEN cDNA 2900041A09 gene		Mm.31025	RIKEN cDNA 2310015K15 gene

ΕP	1	529	838	Α1	

Mm.30412	RIKEN cDNA 5430400P17 gene		Mm.29230	RIKEN cDNA 1500017E18 gene
Mm.30355	(Manual) KIF5A Neuronal Kinesin heavy		Mm.29227	RIKEN cDNA 2300002D11 gene
	chain		Mm.29205	bruno-like 4, RNA binding protein (Droso-
Mm.30266	hemoglobin, beta adult major chain			phila)
Mm.30266	hemoglobin, beta adult major chain	5	Mm.29205	bruno-like 4, RNA binding protein (Droso-
Mm.30206	ATPase, H+ transporting, lysosomal			phila)
M 00450	34kD, V1 subunit D		Mm.2918	megakaryocyte-associated tyrosine ki-
Mm.30156	protease, serine, 11 (Igf binding)		M. 00444	nase
Mm.30155	ATPase, H+ transporting, lysosomal	10	Mm.29141	RIKEN cDNA 0710008N11 gene
Mm.30150	16kD, V0 subunit C RIKEN cDNA 1010001M12 gene	10	Mm.29124 Mm.29075	phosphatidic acid phosphatase type 2B
Mm.30126	membrane interacting protein of RGS16		WIII1.29075	(Manual) Reticulon 1 protein, major inter- nal tag
Mm.30085	aldo-keto reductase family 1, member A4		Mm.29027	SPARC-like 1 (mast9, hevin)
	(aldehyde reductase)		Mm.29027	SPARC-like 1 (mast9, hevin)
Mm.30072	cytochrome c oxidase subunit VIIa poly-	15	Mm.2902	protein tyrosine phosphatase, receptor-
	peptide 2-like			type, N
Mm.30059	myristoylated alanine rich protein kinase		Mm.28955	RIKEN cDNA 4930570C03 gene
	C substrate		Mm.28650	RAB6, member RAS oncogene family
Mm.29976	septin 5		Mm.28650	RAB6, member RAS oncogene family
Mm.29965	RIKEN cDNA 2410104119 gene	20	Mm.28643	vesicle-associated membrane protein 2
Mm.29947	serine/threonine kinase 11		Mm.28561	protein kinase C, zeta
Mm.29939	RIKEN cDNA 1010001N11 gene		Mm.28518	type I transmembrane protein Fn14
Mm.29937	(Manual assignment) polymorphism of		Mm.28357	microtubule-associated protein 1 light
	Mm.29937 ESTs, Weakly similar to pre-			chain 3
14 00004	dicted using Genefinder	25	Mm.2815	RIKEN cDNA 1110021H02 gene
Mm.29921	RAS protein-specific guanine nucleotide-		Mm.28107	ectonucleotide pyrophosphatase/phos-
Mm.2992	releasing factor 1		M 00050	phodiesterase 2
WIIII.2992	(Manual assignment) MBP myelin basic protein		Mm.28058	NADH dehydrogenase (ubiquinone) 1
Mm 00070	•			beta subcomplex 5
		30		DIVEN ADMA 2410011 CO2 cons
Mm.29870 Mm.29867	integral membrane protein 3  NADH dehydrogenase (uhiguipone) 1 al-	30	Mm.27886	RIKEN cDNA 2410011G03 gene
Mm.29867	NADH dehydrogenase (ubiquinone) 1 al-	30		ū
	NADH dehydrogenase (ubiquinone) 1 al- pha subcomplex 2	30	Mm.27886 Mm.27608	Mus musculus, Similar to chromosome
Mm.29867	NADH dehydrogenase (ubiquinone) 1 al- pha subcomplex 2 (Manual) Neurogranin	30		Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC:
Mm.29867 Mm.29857	NADH dehydrogenase (ubiquinone) 1 al- pha subcomplex 2	<i>30</i>		Mus musculus, Similar to chromosome
Mm.29867 Mm.29857	NADH dehydrogenase (ubiquinone) 1 al- pha subcomplex 2 (Manual) Neurogranin Mus musculus, clone IMAGE:5102170,			Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC: 19388 IMAGE:2812475, mRNA, com-
Mm.29867 Mm.29857 Mm.29852	NADH dehydrogenase (ubiquinone) 1 al- pha subcomplex 2 (Manual) Neurogranin Mus musculus, clone IMAGE:5102170, mRNA, partial cds		Mm.27608	Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC: 19388 IMAGE:2812475, mRNA, com- plete cds
Mm.29867 Mm.29857 Mm.29852	NADH dehydrogenase (ubiquinone) 1 al- pha subcomplex 2 (Manual) Neurogranin Mus musculus, clone IMAGE:5102170, mRNA, partial cds Mus musculus, Similar to NDRG family, member 4, clone MGC:7067 IMAGE: 3156802, mRNA, complete cds		Mm.27608 Mm.2755	Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC: 19388 IMAGE:2812475, mRNA, com- plete cds calbindin 2
Mm.29867 Mm.29857 Mm.29852 Mm.29846	NADH dehydrogenase (ubiquinone) 1 al- pha subcomplex 2 (Manual) Neurogranin Mus musculus, clone IMAGE:5102170, mRNA, partial cds Mus musculus, Similar to NDRG family, member 4, clone MGC:7067 IMAGE: 3156802, mRNA, complete cds NADH dehydrogenase flavoprotein 1		Mm.27608 Mm.2755 Mm.27499	Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC: 19388 IMAGE:2812475, mRNA, com- plete cds calbindin 2 RIKEN cDNA 2010004E11 gene RecQ protein-like discs, large homolog 4 (Drosophila)
Mm.29867 Mm.29857 Mm.29852 Mm.29846 Mm.29842 Mm.29823	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex 2 (Manual) Neurogranin Mus musculus, clone IMAGE:5102170, mRNA, partial cds Mus musculus, Similar to NDRG family, member 4, clone MGC:7067 IMAGE: 3156802, mRNA, complete cds NADH dehydrogenase flavoprotein 1 microsomal glutathione S-transferase 3		Mm.27608 Mm.2755 Mm.27499 Mm.27407	Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC: 19388 IMAGE:2812475, mRNA, complete cds calbindin 2 RIKEN cDNA 2010004E11 gene RecQ protein-like discs, large homolog 4 (Drosophila) mitogen activated protein kinase 8 in-
Mm.29867 Mm.29857 Mm.29852 Mm.29846 Mm.29842 Mm.29823 Mm.29807	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex 2 (Manual) Neurogranin Mus musculus, clone IMAGE:5102170, mRNA, partial cds Mus musculus, Similar to NDRG family, member 4, clone MGC:7067 IMAGE: 3156802, mRNA, complete cds NADH dehydrogenase flavoprotein 1 microsomal glutathione S-transferase 3 ubiquitin carboxy-terminal hydrolase L1	35	Mm.27508 Mm.2755 Mm.27499 Mm.27407 Mm.27256 Mm.2720	Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC: 19388 IMAGE:2812475, mRNA, complete cds calbindin 2 RIKEN cDNA 2010004E11 gene RecQ protein-like discs, large homolog 4 (Drosophila) mitogen activated protein kinase 8 interacting protein
Mm.29867 Mm.29857 Mm.29852 Mm.29846 Mm.29842 Mm.29823 Mm.29807 Mm.29807	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex 2 (Manual) Neurogranin Mus musculus, clone IMAGE:5102170, mRNA, partial cds Mus musculus, Similar to NDRG family, member 4, clone MGC:7067 IMAGE: 3156802, mRNA, complete cds NADH dehydrogenase flavoprotein 1 microsomal glutathione S-transferase 3 ubiquitin carboxy-terminal hydrolase L1 ubiquitin carboxy-terminal hydrolase L1	35	Mm.27508 Mm.2755 Mm.27499 Mm.27407 Mm.27256 Mm.2720	Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC: 19388 IMAGE:2812475, mRNA, complete cds calbindin 2 RIKEN cDNA 2010004E11 gene RecQ protein-like discs, large homolog 4 (Drosophila) mitogen activated protein kinase 8 interacting protein RIKEN cDNA 0610043B10 gene
Mm.29867 Mm.29857 Mm.29852 Mm.29846 Mm.29842 Mm.29823 Mm.29807	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex 2 (Manual) Neurogranin Mus musculus, clone IMAGE:5102170, mRNA, partial cds Mus musculus, Similar to NDRG family, member 4, clone MGC:7067 IMAGE: 3156802, mRNA, complete cds NADH dehydrogenase flavoprotein 1 microsomal glutathione S-transferase 3 ubiquitin carboxy-terminal hydrolase L1 ubiquitin carboxy-terminal hydrolase L1 ATPase, H+ transporting, lysosomal	35	Mm.27508 Mm.2755 Mm.27499 Mm.27407 Mm.27256 Mm.2720 Mm.27114 Mm.27087	Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC: 19388 IMAGE:2812475, mRNA, complete cds calbindin 2 RIKEN cDNA 2010004E11 gene RecQ protein-like discs, large homolog 4 (Drosophila) mitogen activated protein kinase 8 interacting protein RIKEN cDNA 0610043B10 gene RIKEN cDNA 2010012C24 gene
Mm.29867 Mm.29857 Mm.29852 Mm.29846 Mm.29842 Mm.29823 Mm.29807 Mm.29807 Mm.29771	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex 2 (Manual) Neurogranin Mus musculus, clone IMAGE:5102170, mRNA, partial cds Mus musculus, Similar to NDRG family, member 4, clone MGC:7067 IMAGE: 3156802, mRNA, complete cds NADH dehydrogenase flavoprotein 1 microsomal glutathione S-transferase 3 ubiquitin carboxy-terminal hydrolase L1 ubiquitin carboxy-terminal hydrolase L1 ATPase, H+ transporting, lysosomal 70kD, V1 subunit A, isoform 1	<i>35</i>	Mm.27508 Mm.2755 Mm.27499 Mm.27407 Mm.27256 Mm.2720 Mm.27114 Mm.27087 Mm.27005	Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC: 19388 IMAGE:2812475, mRNA, complete cds calbindin 2 RIKEN cDNA 2010004E11 gene RecQ protein-like discs, large homolog 4 (Drosophila) mitogen activated protein kinase 8 interacting protein RIKEN cDNA 0610043B10 gene RIKEN cDNA 2010012C24 gene visinin-like 1
Mm.29867 Mm.29857 Mm.29852 Mm.29846 Mm.29842 Mm.29823 Mm.29807 Mm.29807	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex 2 (Manual) Neurogranin Mus musculus, clone IMAGE:5102170, mRNA, partial cds Mus musculus, Similar to NDRG family, member 4, clone MGC:7067 IMAGE: 3156802, mRNA, complete cds NADH dehydrogenase flavoprotein 1 microsomal glutathione S-transferase 3 ubiquitin carboxy-terminal hydrolase L1 ubiquitin carboxy-terminal hydrolase L1 ATPase, H+ transporting, lysosomal 70kD, V1 subunit A, isoform 1 3-monooxgenase/tryptophan 5-monoox-	35	Mm.27508 Mm.2755 Mm.27499 Mm.27407 Mm.27256 Mm.2720 Mm.27114 Mm.27087 Mm.27005 Mm.26633	Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC: 19388 IMAGE:2812475, mRNA, complete cds calbindin 2 RIKEN cDNA 2010004E11 gene RecQ protein-like discs, large homolog 4 (Drosophila) mitogen activated protein kinase 8 interacting protein RIKEN cDNA 0610043B10 gene RIKEN cDNA 2010012C24 gene visinin-like 1 PH domain containing protein in retina 1
Mm.29867 Mm.29857 Mm.29852 Mm.29846 Mm.29842 Mm.29823 Mm.29807 Mm.29807 Mm.29771	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex 2 (Manual) Neurogranin Mus musculus, clone IMAGE:5102170, mRNA, partial cds Mus musculus, Similar to NDRG family, member 4, clone MGC:7067 IMAGE: 3156802, mRNA, complete cds NADH dehydrogenase flavoprotein 1 microsomal glutathione S-transferase 3 ubiquitin carboxy-terminal hydrolase L1 ubiquitin carboxy-terminal hydrolase L1 ATPase, H+ transporting, lysosomal 70kD, V1 subunit A, isoform 1 3-monooxgenase/tryptophan 5-monooxgenase activation protein, gamma poly-	<i>35</i>	Mm.27508  Mm.2755  Mm.27499  Mm.27407  Mm.27256  Mm.2720  Mm.27114  Mm.27087  Mm.27005  Mm.26633  Mm.26633	Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC: 19388 IMAGE:2812475, mRNA, complete cds calbindin 2 RIKEN cDNA 2010004E11 gene RecQ protein-like discs, large homolog 4 (Drosophila) mitogen activated protein kinase 8 interacting protein RIKEN cDNA 0610043B10 gene RIKEN cDNA 2010012C24 gene visinin-like 1 PH domain containing protein in retina 1 PH domain containing protein in retina 1
Mm.29867 Mm.29857 Mm.29852 Mm.29846 Mm.29842 Mm.29823 Mm.29807 Mm.29807 Mm.29711	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex 2 (Manual) Neurogranin Mus musculus, clone IMAGE:5102170, mRNA, partial cds Mus musculus, Similar to NDRG family, member 4, clone MGC:7067 IMAGE: 3156802, mRNA, complete cds NADH dehydrogenase flavoprotein 1 microsomal glutathione S-transferase 3 ubiquitin carboxy-terminal hydrolase L1 ubiquitin carboxy-terminal hydrolase L1 ATPase, H+ transporting, lysosomal 70kD, V1 subunit A, isoform 1 3-monooxgenase/tryptophan 5-monooxgenase activation protein, gamma polypeptide	<i>35</i>	Mm.27508  Mm.2755  Mm.27499  Mm.27407  Mm.27256  Mm.2720  Mm.27114  Mm.27087  Mm.27005  Mm.26633  Mm.26633  Mm.26550	Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC: 19388 IMAGE:2812475, mRNA, complete cds calbindin 2 RIKEN cDNA 2010004E11 gene RecQ protein-like discs, large homolog 4 (Drosophila) mitogen activated protein kinase 8 interacting protein RIKEN cDNA 0610043B10 gene RIKEN cDNA 2010012C24 gene visinin-like 1 PH domain containing protein in retina 1 PH domain containing protein in retina 1 phosphofructokinase, muscle
Mm.29867 Mm.29857 Mm.29852 Mm.29846 Mm.29842 Mm.29823 Mm.29807 Mm.29807 Mm.29711	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex 2 (Manual) Neurogranin Mus musculus, clone IMAGE:5102170, mRNA, partial cds Mus musculus, Similar to NDRG family, member 4, clone MGC:7067 IMAGE: 3156802, mRNA, complete cds NADH dehydrogenase flavoprotein 1 microsomal glutathione S-transferase 3 ubiquitin carboxy-terminal hydrolase L1 ubiquitin carboxy-terminal hydrolase L1 ATPase, H+ transporting, lysosomal 70kD, V1 subunit A, isoform 1 3-monooxgenase/tryptophan 5-monooxgenase activation protein, gamma polypeptide adrenergic receptor kinase, beta 1	<i>35</i>	Mm.27508  Mm.2755  Mm.27499  Mm.27407  Mm.27256  Mm.2720  Mm.27114  Mm.27087  Mm.27005  Mm.26633  Mm.26633	Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC: 19388 IMAGE:2812475, mRNA, complete cds calbindin 2 RIKEN cDNA 2010004E11 gene RecQ protein-like discs, large homolog 4 (Drosophila) mitogen activated protein kinase 8 interacting protein RIKEN cDNA 0610043B10 gene RIKEN cDNA 2010012C24 gene visinin-like 1 PH domain containing protein in retina 1 PH domain containing protein in retina 1 phosphofructokinase, muscle eukaryotic translation elongation factor
Mm.29867 Mm.29857 Mm.29852 Mm.29846 Mm.29842 Mm.29823 Mm.29807 Mm.29807 Mm.29711	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex 2 (Manual) Neurogranin Mus musculus, clone IMAGE:5102170, mRNA, partial cds Mus musculus, Similar to NDRG family, member 4, clone MGC:7067 IMAGE: 3156802, mRNA, complete cds NADH dehydrogenase flavoprotein 1 microsomal glutathione S-transferase 3 ubiquitin carboxy-terminal hydrolase L1 ubiquitin carboxy-terminal hydrolase L1 ATPase, H+ transporting, lysosomal 70kD, V1 subunit A, isoform 1 3-monooxgenase/tryptophan 5-monooxgenase activation protein, gamma polypeptide adrenergic receptor kinase, beta 1 actin, beta, cytoplasmic	<i>35</i>	Mm.27508  Mm.2755  Mm.27499  Mm.27407  Mm.27256  Mm.2720  Mm.27114  Mm.27087  Mm.27005  Mm.26633  Mm.26633  Mm.26550  Mm.2645	Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC: 19388 IMAGE:2812475, mRNA, complete cds calbindin 2 RIKEN cDNA 2010004E11 gene RecQ protein-like discs, large homolog 4 (Drosophila) mitogen activated protein kinase 8 interacting protein RIKEN cDNA 0610043B10 gene RIKEN cDNA 2010012C24 gene visinin-like 1 PH domain containing protein in retina 1 phosphofructokinase, muscle eukaryotic translation elongation factor 1 alpha 2
Mm.29867 Mm.29857 Mm.29852 Mm.29846 Mm.29842 Mm.29823 Mm.29807 Mm.29711 Mm.29711 Mm.29711	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex 2 (Manual) Neurogranin Mus musculus, clone IMAGE:5102170, mRNA, partial cds Mus musculus, Similar to NDRG family, member 4, clone MGC:7067 IMAGE: 3156802, mRNA, complete cds NADH dehydrogenase flavoprotein 1 microsomal glutathione S-transferase 3 ubiquitin carboxy-terminal hydrolase L1 ubiquitin carboxy-terminal hydrolase L1 ATPase, H+ transporting, lysosomal 70kD, V1 subunit A, isoform 1 3-monooxgenase/tryptophan 5-monooxgenase activation protein, gamma polypeptide adrenergic receptor kinase, beta 1	35 40 45	Mm.27508  Mm.2755  Mm.27499  Mm.27407  Mm.27256  Mm.2720  Mm.27114  Mm.27087  Mm.27005  Mm.26633  Mm.26633  Mm.26550	Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC: 19388 IMAGE:2812475, mRNA, complete cds calbindin 2 RIKEN cDNA 2010004E11 gene RecQ protein-like discs, large homolog 4 (Drosophila) mitogen activated protein kinase 8 interacting protein RIKEN cDNA 0610043B10 gene RIKEN cDNA 2010012C24 gene visinin-like 1 PH domain containing protein in retina 1 PH domain containing protein in retina 1 phosphofructokinase, muscle eukaryotic translation elongation factor
Mm.29867 Mm.29852 Mm.29846 Mm.29846 Mm.29842 Mm.29823 Mm.29807 Mm.29711 Mm.29711 Mm.297 Mm.29633	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex 2 (Manual) Neurogranin Mus musculus, clone IMAGE:5102170, mRNA, partial cds Mus musculus, Similar to NDRG family, member 4, clone MGC:7067 IMAGE: 3156802, mRNA, complete cds NADH dehydrogenase flavoprotein 1 microsomal glutathione S-transferase 3 ubiquitin carboxy-terminal hydrolase L1 ubiquitin carboxy-terminal hydrolase L1 ATPase, H+ transporting, lysosomal 70kD, V1 subunit A, isoform 1 3-monooxgenase/tryptophan 5-monooxgenase activation protein, gamma polypeptide adrenergic receptor kinase, beta 1 actin, beta, cytoplasmic RIKEN cDNA 1810008021 gene	35 40 45	Mm.27608  Mm.2755  Mm.27499  Mm.27407  Mm.27256  Mm.2720  Mm.27114  Mm.27087  Mm.27005  Mm.26633  Mm.26633  Mm.26550  Mm.2645  Mm.2635	Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC: 19388 IMAGE:2812475, mRNA, complete cds calbindin 2 RIKEN cDNA 2010004E11 gene RecQ protein-like discs, large homolog 4 (Drosophila) mitogen activated protein kinase 8 interacting protein RIKEN cDNA 0610043B10 gene RIKEN cDNA 2010012C24 gene visinin-like 1 PH domain containing protein in retina 1 PH domain containing protein in retina 1 phosphofructokinase, muscle eukaryotic translation elongation factor 1 alpha 2 pyruvate kinase 3 cholecystokinin
Mm.29867 Mm.29852 Mm.29846 Mm.29846 Mm.29842 Mm.29823 Mm.29807 Mm.29711 Mm.29711 Mm.297 Mm.29633	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex 2 (Manual) Neurogranin Mus musculus, clone IMAGE:5102170, mRNA, partial cds Mus musculus, Similar to NDRG family, member 4, clone MGC:7067 IMAGE: 3156802, mRNA, complete cds NADH dehydrogenase flavoprotein 1 microsomal glutathione S-transferase 3 ubiquitin carboxy-terminal hydrolase L1 ubiquitin carboxy-terminal hydrolase L1 ATPase, H+ transporting, lysosomal 70kD, V1 subunit A, isoform 1 3-monooxgenase/tryptophan 5-monooxgenase activation protein, gamma polypeptide adrenergic receptor kinase, beta 1 actin, beta, cytoplasmic RIKEN cDNA 1810008021 gene Mus musculus, clone IMAGE:3964267,	35 40 45	Mm.27608  Mm.2755  Mm.27499  Mm.27407  Mm.27256  Mm.2720  Mm.27114  Mm.27087  Mm.27005  Mm.26633  Mm.26633  Mm.26550  Mm.2645  Mm.2635  Mm.2635  Mm.2635	Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC: 19388 IMAGE:2812475, mRNA, complete cds calbindin 2 RIKEN cDNA 2010004E11 gene RecQ protein-like discs, large homolog 4 (Drosophila) mitogen activated protein kinase 8 interacting protein RIKEN cDNA 0610043B10 gene RIKEN cDNA 2010012C24 gene visinin-like 1 PH domain containing protein in retina 1 PH domain containing protein in retina 1 phosphofructokinase, muscle eukaryotic translation elongation factor 1 alpha 2 pyruvate kinase 3
Mm.29867 Mm.29857 Mm.29852 Mm.29846 Mm.29842 Mm.29823 Mm.29807 Mm.29711 Mm.29717 Mm.29711 Mm.297 Mm.29633 Mm.29600	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex 2 (Manual) Neurogranin Mus musculus, clone IMAGE:5102170, mRNA, partial cds Mus musculus, Similar to NDRG family, member 4, clone MGC:7067 IMAGE: 3156802, mRNA, complete cds NADH dehydrogenase flavoprotein 1 microsomal glutathione S-transferase 3 ubiquitin carboxy-terminal hydrolase L1 ubiquitin carboxy-terminal hydrolase L1 ATPase, H+ transporting, lysosomal 70kD, V1 subunit A, isoform 1 3-monooxgenase/tryptophan 5-monooxgenase activation protein, gamma polypeptide adrenergic receptor kinase, beta 1 actin, beta, cytoplasmic RIKEN cDNA 1810008021 gene Mus musculus, clone IMAGE:3964267, mRNA	35 40 45	Mm.27608  Mm.2755  Mm.27499  Mm.27407  Mm.27256  Mm.2720  Mm.27114  Mm.27087  Mm.27005  Mm.26633  Mm.26633  Mm.26550  Mm.2645  Mm.2635  Mm.2635  Mm.2635  Mm.2635	Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC: 19388 IMAGE:2812475, mRNA, complete cds calbindin 2 RIKEN cDNA 2010004E11 gene RecQ protein-like discs, large homolog 4 (Drosophila) mitogen activated protein kinase 8 interacting protein RIKEN cDNA 0610043B10 gene RIKEN cDNA 2010012C24 gene visinin-like 1 PH domain containing protein in retina 1 PH domain containing protein in retina 1 phosphofructokinase, muscle eukaryotic translation elongation factor 1 alpha 2 pyruvate kinase 3 cholecystokinin RIKEN cDNA 2010003014 gene RIKEN cDNA 2900002P20 gene ring finger protein 11
Mm.29867 Mm.29852 Mm.29846 Mm.29846 Mm.29842 Mm.29823 Mm.29807 Mm.29711 Mm.29717 Mm.29711 Mm.297 Mm.29633 Mm.29600 Mm.2948	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex 2 (Manual) Neurogranin Mus musculus, clone IMAGE:5102170, mRNA, partial cds Mus musculus, Similar to NDRG family, member 4, clone MGC:7067 IMAGE: 3156802, mRNA, complete cds NADH dehydrogenase flavoprotein 1 microsomal glutathione S-transferase 3 ubiquitin carboxy-terminal hydrolase L1 ubiquitin carboxy-terminal hydrolase L1 ATPase, H+ transporting, lysosomal 70kD, V1 subunit A, isoform 1 3-monooxgenase/tryptophan 5-monooxgenase activation protein, gamma polypeptide adrenergic receptor kinase, beta 1 actin, beta, cytoplasmic RIKEN cDNA 1810008021 gene Mus musculus, clone IMAGE:3964267, mRNA H2-K region expressed gene 2 SCAN domain-containing 1 RIKEN cDNA 1810011001 gene	35 40 45	Mm.27608  Mm.2755  Mm.27499  Mm.27407  Mm.27256  Mm.2720  Mm.27114  Mm.27087  Mm.27005  Mm.26633  Mm.26633  Mm.26550  Mm.2645  Mm.2635  Mm.2635  Mm.2635  Mm.2635  Mm.2635  Mm.2635	Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC: 19388 IMAGE:2812475, mRNA, complete cds calbindin 2 RIKEN cDNA 2010004E11 gene RecQ protein-like discs, large homolog 4 (Drosophila) mitogen activated protein kinase 8 interacting protein RIKEN cDNA 0610043B10 gene RIKEN cDNA 2010012C24 gene visinin-like 1 PH domain containing protein in retina 1 PH domain containing protein in retina 1 phosphofructokinase, muscle eukaryotic translation elongation factor 1 alpha 2 pyruvate kinase 3 cholecystokinin RIKEN cDNA 2010003014 gene RIKEN cDNA 2900002P20 gene ring finger protein 11 NCK-associated protein 1
Mm.29867 Mm.29857 Mm.29852 Mm.29846 Mm.29842 Mm.29823 Mm.29807 Mm.29807 Mm.29771 Mm.29717 Mm.29711 Mm.297 Mm.29633 Mm.29600 Mm.2948 Mm.2948 Mm.29477 Mm.29415 Mm.29362	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex 2 (Manual) Neurogranin Mus musculus, clone IMAGE:5102170, mRNA, partial cds Mus musculus, Similar to NDRG family, member 4, clone MGC:7067 IMAGE: 3156802, mRNA, complete cds NADH dehydrogenase flavoprotein 1 microsomal glutathione S-transferase 3 ubiquitin carboxy-terminal hydrolase L1 ubiquitin carboxy-terminal hydrolase L1 ATPase, H+ transporting, lysosomal 70kD, V1 subunit A, isoform 1 3-monooxgenase/tryptophan 5-monooxgenase activation protein, gamma polypeptide adrenergic receptor kinase, beta 1 actin, beta, cytoplasmic RIKEN cDNA 1810008021 gene Mus musculus, clone IMAGE:3964267, mRNA H2-K region expressed gene 2 SCAN domain-containing 1 RIKEN cDNA 1810011001 gene expressed sequence Al414999	35 40 45	Mm.27608  Mm.2755  Mm.27499  Mm.27407  Mm.27256  Mm.2720  Mm.27114  Mm.27087  Mm.27005  Mm.26633  Mm.26633  Mm.26550  Mm.2645  Mm.2635	Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC: 19388 IMAGE:2812475, mRNA, complete cds calbindin 2 RIKEN cDNA 2010004E11 gene RecQ protein-like discs, large homolog 4 (Drosophila) mitogen activated protein kinase 8 interacting protein RIKEN cDNA 0610043B10 gene RIKEN cDNA 2010012C24 gene visinin-like 1 PH domain containing protein in retina 1 PH domain containing protein in retina 1 phosphofructokinase, muscle eukaryotic translation elongation factor 1 alpha 2 pyruvate kinase 3 cholecystokinin RIKEN cDNA 2010003014 gene RIKEN cDNA 2900002P20 gene ring finger protein 11 NCK-associated protein 1 internexin neuronal intermediate fila-
Mm.29867 Mm.29857 Mm.29852 Mm.29846 Mm.29842 Mm.29823 Mm.29807 Mm.29807 Mm.29771 Mm.29717 Mm.29711 Mm.297 Mm.29633 Mm.29600 Mm.2948 Mm.29477 Mm.29415	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex 2 (Manual) Neurogranin Mus musculus, clone IMAGE:5102170, mRNA, partial cds Mus musculus, Similar to NDRG family, member 4, clone MGC:7067 IMAGE: 3156802, mRNA, complete cds NADH dehydrogenase flavoprotein 1 microsomal glutathione S-transferase 3 ubiquitin carboxy-terminal hydrolase L1 ubiquitin carboxy-terminal hydrolase L1 ATPase, H+ transporting, lysosomal 70kD, V1 subunit A, isoform 1 3-monooxgenase/tryptophan 5-monooxgenase activation protein, gamma polypeptide adrenergic receptor kinase, beta 1 actin, beta, cytoplasmic RIKEN cDNA 1810008021 gene Mus musculus, clone IMAGE:3964267, mRNA H2-K region expressed gene 2 SCAN domain-containing 1 RIKEN cDNA 1810011001 gene	35 40 45	Mm.27608  Mm.2755  Mm.27499  Mm.27407  Mm.27256  Mm.2720  Mm.27114  Mm.27087  Mm.27005  Mm.26633  Mm.26633  Mm.26550  Mm.2645  Mm.2635  Mm.2635	Mus musculus, Similar to chromosome 9 open reading frame 16, clone MGC: 19388 IMAGE:2812475, mRNA, complete cds calbindin 2 RIKEN cDNA 2010004E11 gene RecQ protein-like discs, large homolog 4 (Drosophila) mitogen activated protein kinase 8 interacting protein RIKEN cDNA 0610043B10 gene RIKEN cDNA 2010012C24 gene visinin-like 1 PH domain containing protein in retina 1 PH domain containing protein in retina 1 phosphofructokinase, muscle eukaryotic translation elongation factor 1 alpha 2 pyruvate kinase 3 cholecystokinin RIKEN cDNA 2010003014 gene RIKEN cDNA 2900002P20 gene ring finger protein 11 NCK-associated protein 1

E	P 1	529	838 A1	ļ
---	-----	-----	--------	---

Mm.2446	synaptotagmin 4			pha o
Mm.24376	Mus musculus mRNA for calsyntenin-3 (Cs3 gene)		Mm.20964	guanine nucleotide binding protein, alpha o
Mm.2411	Ras-GTPase-activating protein (GAP		Mm.2082	apolipoprotein D
	<120>) SH3-domain binding protein 2	5	Mm.206218	Mus musculus, Similar to hypothetical
Mm.24092	N-ethylmaleimide sensitive fusion protein			protein FLJ22237, clone MGC:27683 IMAGE:4913322, mRNA, complete cds
Mm.24092	N-ethylmaleimide sensitive fusion protein		Mm.2060 Mm.20472	RIKEN cDNA 2900010105 gene vertebrate homolog of C. elegans Lin-7
Mm.2400	glutathione peroxidase 4	10		type 2
Mm.2397	synaptophysin		Mm.203939	expressed sequence Al256814
Mm.23826	phosphotyrosyl phosphatase activator		Mm.203924	expressed sequence AW259572
Mm.2381	amyloid beta (A4) precursor-like protein		Mm.203921	expressed sequence Al850305
	1		Mm.202728	expressed sequence Al447901
Mm.2338	glutamine synthetase	15	Mm.202696	expressed sequence AA409221
Mm.2338	glutamine synthetase		Mm.201729	expressed sequence Al426007
Mm.2326	macrophage migration inhibitory factor		Mm.2011	glutathione S-transferase, mu 1
Mm.2319	Scgn10 like-protein		Mm.200858	RIKEN cDNA 2410129E14 gene
Mm.23023	RIKEN cDNA 1500009C09 gene		Mm.200843	synuclein, beta
Mm.23002	RIKEN cDNA 5330410G16 gene	20	Mm.200817	expressed sequence AW124717
Mm.22699	selenoprotein P, plasma, 1		Mm.200817	expressed sequence AW124717
Mm.22637	RIKEN cDNA 0910001L24 gene		Mm.200806	(Manual) no clear assignment, probably
Mm.22597	RIKEN cDNA 2310042E05 gene			non-coding (but spliced) RNA gene
Mm.22473	Rab acceptor 1 (prenylated)		Mm.200511	expressed sequence Al115024
Mm.22149	succinate dehydrogenase complex,	25	Mm.199903	expressed sequence Al850290
	subunit A, flavoprotein (Fp)		Mm.199652	expressed sequence Al838505
Mm.2214	septin 4		Mm.198588	expressed sequence Al851970
Mm.220966	reticulon 4		Mm.19834	RIKEN cDNA 0610033L03 gene
Mm.220898	calmodulin 3		Mm.197523	brain acyl-CoA hydrolase
Mm.220885	neurochondrin	30	Mm.196614	eukaryotic translation elongation factor
Mm.2206	NADH dehydrogenase (ubiquinone) fla-			1 alpha 1
	voprotein 2		Mm.196611	synapsin I
Mm.219776	RIKEN cDNA 1110001E17 gene		Mm.196607	eukaryotic translation initiation factor 5A
Mm.218848	RIKEN cDNA 3010002G01 gene		Mm.196605	hexokinase 1
Mm.218764	guanine nucleotide binding protein 13,	35	Mm.196578	mitochondrial carrier homolog 1
	gamma		Mm.196344	lusterin
Mm.218611	receptor (calcitonin) activity modifying		Mm.196239	RIKEN cDNA 4922501H04 gene
M. 04740	protein 2		Mm.195869	ATPase, H+ transporting, lysosomal
Mm.21743	malate dehydrogenase, mitochondrial	40	M 4050	31kDa, V1 subunit E
Mm.216438	Mus musculus, clone IMAGE:5068657,	40	Mm.1956	neurofilament, light polypeptide
M 01 00 40	mRNA, partial cds		Mm.19370	ATP synthase, H+ transporting, mi-
Mm.216240	Mus musculus, clone IMAGE:3594799,		Mm 102520	tochondrial F1F0 complex, subunit e
Mm 01405	mRNA RIKEN cDNA 2610102M01 gene		Mm.193539	H1 histone family, member 2
Mm.21485 Mm.214549	3	45	Mm.192991	Mus musculus, Similar to metallot-
WIII.214549	Mus musculus, Similar to vesicle-associated calmodulin-binding protein, clone	43		hionein 1, clone MGC:27821 IMAGE: 3483861, mRNA, complete cds
	MGC:28873 IMAGE:4527857, mRNA,		Mm.19133	amyloid beta (A4) precursor-like protein
	complete cds		WIIII. 13133	2
Mm.2133	centaurin, gamma 3		Mm.19047	expressed sequence Al425998
Mm.212672	S100 protein, beta polypeptide, neural	50	Mm.182912	growth hormone inducible transmem-
Mm.212516	RIKEN cDNA 2900002L20 gene	00	WIII. 102312	brane protein
Mm.21251	deleted in polyposis 1		Mm.18218	ganglioside-induced differentiation-as-
Mm.21162	genes associated with retinoid-IFN-in-			sociated-protein 1
	duced mortality 19		Mm.181894	RIKEN cDNA 2900092E17 gene
Mm.2108	transthyretin	55	Mm.181721	RIKEN cDNA 2610041P16 gene
Mm.21071	ADP-ribosylation-like 2		Mm.180182	cytochrome c oxidase, subunit Vb
Mm.21069	RIKEN cDNA 0610007A03 gene		Mm.1776	ferritin heavy chain
Mm.20964	guanine nucleotide binding protein, al-		Mm.177272	brain protein 17
				•

17	EP 1 529 838 A1

Mm.177117	Mus musculus, clone MGC:31632 IMAGE:4511454, mRNA, complete cds		Mm.115124 Mm.114810	brain protein 14 expressed sequence AW060990
Mm.176927	RIKEN cDNA 2610301115 gene		Mm.1147	Mus musculus calmodulin III (Calm3)
Mm.17484	synuclein, alpha	_	M 40707	mRNA, 3' untranslated region
Mm.16831	creatine kinase, brain	5	Mm.10727	ATPase, H+ transporting, lysosomal
Mm.16769	RIKEN cDNA 5031406P05 gene			56/58kD, V1 subunit B, isoform 2
Mm.16767	heterogeneous nuclear ribonucleoprotein A2/B1		Mm.103709	potassium inwardly-rectifying channel, subfamily J, member 10
Mm.16763	aldolase 1, A isoform		Mm.103605	DnaJ (Hsp40) homolog, subfamily B,
Mm.16228	solute carrier family 25 (mitochondrial	10		member 10
	carrier; adenine nucleotide transloca-		Mm.102278	secretory carrier membrane protein 5
	tor), member 4		Mm.102244	expressed sequence R74975
Mm.16080	dynamin		Mm.101476	(Manual assignment) BNPI, VGLUT-1,
Mm.158871	RIKEN cDNA 2410003L22 gene			mouse homolog of putative vesicular
Mm.157929	ESTs, Weakly similar to PBAS MOUSE PROBASIN PRECURSOR	15		glutamate transporter, Na+/Phosphate cotransporter
Mm.157859	ESTs		Mm.100980	calneuron 1
Mm.157648	RIKEN cDNA 5730403B10 gene		Mm.1008	prostaglandin D2 synthase (21 kDa,
Mm.15711	cyclic nucleotide phosphodiesterase 1			brain)
Mm.156959	beta-spectrin 4	20	Mm.1008	(Manual) Prostaglandin H2 D-Isomera-
Mm.15571	amyloid beta (A4) precursor protein			se (PGD2 SYNTHASE) (PGDS2)
Mm.15512	potassium voltage-gated channel, sha-			(PGDS) member of lipocalin family
	ker-related subfamily, beta member 2			( and ) manual of apodam raining
Mm.154651	purine rich element binding protein B		Liste D: Posit	tivmarker neurale Stammzellen (1.);
Mm.153758	RIKEN cDNA 0610040H15 gene	25		Total otal in the control of the con
Mm.15125	stromal cell derived factor receptor 1		ES-Zeilen +· I	PSA-NCAM - ; Adult brain -
Mm.14798	ribosomal protein S13			, naut blain
Mm.142511	expressed sequence Al173355		[0042]	
Mm.142187	RIKEN cDNA 2610009E16 gene		[00.2]	
Mm.142140	neurofilament, medium polypeptide	30	Mm.9703	(Manual) copper transport protein/cha-
Mm.140761	DnaJ (Hsp40) homolog, subfamily C,			perone ATOX1
	member 5		Mm.930	cathepsin L
Mm.139797	expressed sequence Al848587		Mm.90787	nerve growth factor receptor
Mm. 139239	RIKEN cDNA 2900016C05 gene		,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	(TNFRSF16) associated protein 1
Mm.139239	RIKEN cDNA 2900016C05 gene	35	Mm.90587	enolase 1, alpha non-neuron
Mm.139239	RIKEN cDNA 2900016C05 gene		Mm.90115	lysophospholipase 1
Mm.138866	apolipoprotein E		Mm.90003	gap junction membrane channel prote-
Mm.13859	ribosomal protein L41			in beta 3
Mm.1383	Rho GDP dissociation inhibitor (GDI)		Mm.88302	EST, Weakly similar to S14234 hypo-
	gamma	40		thetical protein - mouse
Mm.135621	expressed sequence Al848120		Mm.88212	tubulin, alpha 6
Mm.13445	3-oxoacid CoA transferase		Mm.87581	(Manual) fibronectin 1, internal tag (ma-
Mm.1339	chromogranin B			jor tag probably AAAAAAAAAA)
Mm.131127	RIKEN cDNA 6230410L23 gene		Mm.87293	WD repeat domain 12
Mm.12958	kinesin light chain 2	45	Mm.87216	Rab geranylgeranyl transferase, a sub-
Mm.12860	G protein-coupled receptor 37-like 1			unit
Mm.1268	proteolipid protein (myelin)		Mm.8155	TG interacting factor
Mm.1268	(Manual assignment) PLP Myelin Pro-		Mm.78861	nucleolar and coiled-body phosphopro-
	teolipid Protein, uh05d10.r1 Soares			tein 1
	mouse hypothalamus NMHy Mus mus-	50	Mm.76780	ESTs
	culus cDNA clone 1617043 5' similar to		Mm.7417	cyclin D3
	gb:M54927 MYELIN PROTEOLIPID		Mm.7387	RNA polymerase 1-4 (194 kDa subunit)
	PROTEIN		Mm.7381	hypoxia induced gene 1
Mm.12468	glioblastoma amplified sequence		Mm.725	ribosomal protein L7a
Mm.124592	expressed sequence AW214631	55	Mm.71046	ESTs
Mm.1239	glial fibrillary acidic protein		Mm.70127	ribosomal protein L12
Mm.1222	brain abundant, membrane attached si-		Mm.69647	pancreas specific transcription factor,
	gnal protein 2			1a

19

Mm.69049	cDNA sequence AF155546			IMAGE:3992883, mRNA, complete cds
Mm.6700	eukaryotic translation initiation factor		Mm.3845	Mus musculus, eukaryotic translation
	4E binding protein 1			termination factor 1, clone MGC:18745
Mm.66	ribosomal protein S4, X-linked	_		IMAGE:3992883, mRNA, complete cds
Mm.6579	centromere autoantigen A	5	Mm.38151	adenylosuccinate lyase
Mm.6534	calpain, small subunit 1		Mm.38057	ESTs
Mm.6343	nucleophosmin 1		Mm.3776	Mus musculus, clone MGC:37810
Mm.584	annexin A2			IMAGE:5098241, mRNA, complete cds
Mm.57223	helicase, lymphoid specific		Mm.3752	RAN binding protein 1
Mm.57153	sterol O-acyltransferase 2	10	Mm.36241	B-cell receptor-associated protein 37
Mm.5624	DEAD/H (Asp-Glu-Ala-Asp/His) box		Mm.360	cytochrome c oxidase, subunit Va
	polypeptide 16		Mm.3572	RIKEN cDNA 1110033J19 gene
Mm.548	cytochrome c oxidase, subunit VIc		Mm.35621	ESTs
Mm.5305	(Manual) GNB2L1, RACK1, Receptor	45	Mm.35605	cadherin 1
	of activated C kinase, WD40-repeat	15	Mm.3487	ribosomal protein L30
	protein		Mm.3486	ribosomal protein L3
Mm.5290	(Manual) 60S ribosomal protein L17		Mm.34828	heat shock protein, 105 kDa
	(L23) (popey3-annotation wrong)		Mm.34797	cellular retinoic acid binding protein I
Mm.4993	matrix metalloproteinase 3	00	Mm.34606	RIKEN cDNA 2610511F02 gene
Mm.493	CCCTC-binding factor	20	Mm.34554	Mus musculus, Similar to E2F trans-
Mm.4890	Finkel-Biskis-Reilly murine sarcoma vi-			cription factor 4, p107/p130-binding,
	rus (FBR-MuSV) ubiquitously expres-			clone MGC:37558 IMAGE:4987691,
	sed (fox derived)		14 . 0400	mRNA, complete cds
Mm.4770	frizzled homolog 7 (Drosophila)		Mm.3438	lamin A
Mm.4742	proliferation-associated 2G4, 38kD	25	Mm.34351	Mus musculus, Similar to hypothetical
Mm.46461	L-threonine dehydrogenase			protein FLJ13187, clone MGC:28979
Mm.4606	branched chain aminotransferase 1,			IMAGE:4503757, mRNA, complete cds
14 4500	cytosolic		Mm.34102	ornithine decarboxylase, structural
Mm.4560	low density lipoprotein receptor-related		Mm.3379	serine hydroxymethyl transferase 1
	protein associated protein 1	30	N 00040	(soluble)
Mm.45237	RIKEN cDNA 2610318N02 gene		Mm.33240	epithelial V-like antigen
Mm.45151	RIKEN cDNA 1700043E15 gene		Mm.33202	RIKEN cDNA 2410018A17 gene
Mm.4502	mini chromosome maintenance de-		Mm.32879	testis expressed gene 17
N 4000t	ficient (S. cerevisiae)	<i>3</i> 5	Mm.321	secreted phosphoprotein 1
Mm.43831	lectin, galactose binding, soluble 1	33	Mm.318	RIKEN cDNA 2010107E04 gene
Mm.43162	RIKEN cDNA 0710008D09 gene		Mm.31227	expressed sequence AW123847
Mm.42960	RIKEN cDNA 2610301D06 gene		Mm.30929	peroxiredoxin 1
Mm.4280	RIKEN cDNA 2010203J19 gene		Mm.3049	CDC28 protein kinase 1 peptidylprolyl isomerase D (cyclophilin
Mm.42790	ribosomal protein S18	40	Mm.30242	
Mm.42767	ribosomal protein S17	40	Mm 20194	D) RIKEN cDNA 2700086123 gene
Mm.42197	proteasome (prosome, macropain)		Mm.30184 Mm.30114	amyotrophic lateral sclerosis 2 (juveni-
Mm.42196	subunit, beta type 1		WIII1.50114	le) homolog (human)
	nuclear protein 95		Mm.30060	RIKEN cDNA 2310008N12 gene
Mm.42195	RuvB-like protein 1  Mus musculus, clone MGC:28892	45	Mm.30049	complement component 1, q subcom-
Mm.41467	•	45	WIII.30049	ponent binding protein
Mm.41151	IMAGE:4912251, mRNA, complete cds ESTs		Mm.30034	translocase of inner mitochondrial
Mm.41061			141111.30054	membrane 8 homolog a (yeast)
Mm.41	RIKEN cDNA 1810055P05 gene (Manual) Mitochondrial ATP synthase		Mm.29904	mitochondrial ribosomal protein L15
WITH,41	oligomycin sensitivity conferral protein	50	Mm.29902	Mus musculus, Similar to phosphoseri-
	(OSCP) (ATP5O)	55	WIII1.29902	ne aminotransferase, clone MGC:6462
Mm.4095				IMAGE:2616298, mRNA, complete cds
Mm.4095 Mm.4024	inactive X specific transcripts cofilin 1, non-muscle		Mm.29859	eukaryotic translation initiation factor 2,
Mm.3925	S100 calcium binding protein A4			subunit 2 (beta, 38kDa)
Mm.38718	ESTs, Moderately similar to S12207 hy-	55	Mm.29856	RIKEN cDNA 9130022B02 gene
141111.007 10	pothetical protein		Mm.29717	3-monooxgenase/tryptophan 5-mo-
Mm.3845	Mus musculus, eukaryotic translation			nooxgenase activation protein, gamma
	termination factor 1, clone MGC: 18745			polypeptide
	toation factor 1, cione in Go. 10745			F ) Popular

21

Mm.29714	(Manual) mouse version of muscle-			tor), member 13
	specific protein M9		Mm.24506	Mus musculus, clone IMAGE:3591061,
Mm.29675	thioredoxin-like 2			mRNA, partial cds
Mm.29619	RIKEN cDNA 1200007E24 gene		Mm.2437	BING4 protein
Mm.29513	NADH dehydrogenase (ubiquinone) 1	5	Mm.2424	ribosomal protein L10A
	alpha subcomplex, 7 (14.5kD, B14.5a)		Mm.24220	RIKEN cDNA 2310003F16 gene
Mm.29504	sperm specific antigen 1		Mm.24219	RIKEN cDNA 1810037117 gene
Mm.2942	asparagine synthetase		Mm.24174	Mus musculus, similar to alanyl-tRNA
Mm.29405	ring-box 1			synthetase (H. sapiens), clone MGC:
Mm.29363	RIKEN cDNA 2310044F10 gene	10		37368 IMAGE:4976684, mRNA, com-
Mm.2930	Mus musculus, Similar to peter pan			plete cds
	(Drosophila) homolog, clone MGC:		Mm.2395	male enhanced antigen 1
	25669 IMAGE:4489442, mRNA, com-		Mm.2355	prohibitin
Mm 20100	plete cds		Mm.235	ubiquitin B
Mm.29192 Mm.29148	asparaginyl-tRNA synthetase	15	141111.22701	integrin beta 4 binding protein
Mm.29122	RIKEN cDNA 2400008B06 gene		Mm.22626	Mus musculus, Similar to chromosome
Mm.29076	RIKEN cDNA 0610012D09 gene RIKEN cDNA 2510010F10 gene			14 open reading frame 3, clone MGC:
Mm.28919	destrin			36589 IMAGE:5320590, mRNA, com-
Mm.28892	expressed sequence AA959742	20	Mm.2246	plete cds
Mm.28805	SET translocation	20	14111.2240	proteasome (prosome, macropain) subunit, beta type 7
Mm.2849	heat shock protein, 74 kDa, A		Mm.22421	telomerase binding protein, p23
Mm.28483	Mus musculus, Similar to hypothetical		Mm.22421	telomerase binding protein, p23
	protein FLJ22479, clone IMAGE:		Mm.22317	RIKEN cDNA 8430410A17 gene
	4487274, mRNA, partial cds	25		cyclin D1
Mm.28405	serum/glucocorticoid regulated kinase		Mm.22271	smt3-specific isopeptidase 1
Mm.28173	ESTs, Moderately similar to JC5224		Mm.220992	Mus musculus, clone IMAGE:3492506,
	methioninetRNA ligase			mRNA, partial cds
Mm.28053	RIKEN cDNA 1110017C15 gene		Mm.219671	Mus musculus, clone MGC:36369
Mm.28035	ESTs, Weakly similar to	30		IMAGE:4982239, mRNA, complete cds
	TRHY_HUMAN TRICHOHYALI		Mm.219458	RNA binding protein gene with multiple
Mm.27901	RIKEN cDNA 1110020J08 gene		`	splicing
Mm.27858	RIKEN cDNA 1110036B12 gene		Mm.218533	RIKEN cDNA 1500016H10 gene
Mm.27855	replication factor C (activator 1) 2		Mm.2180	heat shock protein, 84 kDa 1
Mm.2758	(40kD)	35	Mm.21758	cytochrome P450, 2e1, ethanol induci-
Mm.27536	makorin, ring finger protein, 3 ESTs, Highly similar to hypothetical		M 04000	ble
WIII.27550	protein FLJ14075		Mm.21630	expressed sequence AU022237
Mm.27526	(Manual) Arginyl tRNA synthetase (RI-		Mm.21569 Mm.213020	RIKEN cDNA 2700069E09 gene
141111.27320	KEN cDNA 2610011N19)	40	WIIII.2 13020	(Manual) 60S ribosomal protein L32 (RPL32)
Mm.27186	Mus musculus, Similar to CG7083 ge-		Mm.212899	Mus musculus, Similar to RIKEN cDNA
	ne product, clone MGC:6480 IMAGE:			1200009K13 gene, clone MGC: 18794
	2646515, mRNA, complete cds			IMAGE:4193513, mRNA, complete cds
Mm.2718	eukaryotic translation elongation factor		Mm.21289	ribosomal protein S12
	1 beta 2	45	Mm.21086	eukaryotic translation elongation factor
Mm.2718	eukaryotic translation elongation factor			1 delta (guanine nucleotide exchange
	1 beta 2			protein)
Mm.27134	RIKEN cDNA 2610033C09 gene		Mm.210638	EST
Mm.265	ribosomal protein S25		Mm.21062	expressed sequence C87860
Mm.2647	profilin 1	50	Mm.21054	nuclease sensitive element binding
Mm.2623	serine (or cysteine) proteinase inhibitor,			protein 1
M 05040	clade B (ovalbumin), member 6		Mm.20943	FK506 binding protein 9
Mm.25642	RIKEN cDNA 2310034K10 gene		Mm.20925	G1 to phase transition 1
Mm.254	tumor protein, translationally-controlled	g e	Mm.20918	nuclear localization signal protein ab-
Mm.25328	1 ESTs	55	Mm 20040	sent in velo-cardio-facial patients
Mm.24513	solute carrier family 25 (mitochondrial		Mm.20848	regulatory factor X-associated ankyrin-
	carrier; adenine nucleotide transloca-		Mm.20847	containing protein sorting nexin 5
			WHITE COM /	Sorting Hexili 3

23	EP 1 529 838 A1

Mm.20290expressed sequence AW536573Mm.154915ribosomalMm.20288glutathione reductase 1Mm.154387transketolMm.200920ribosomal protein S28Mm.153963CD8 antig	DNA 2610034E13 gene I protein S29 Iase gen, beta chain in subunit 6a (zeta)
Mm. 197551heat shock 70kD protein 8Mm.148973RIKEN cDMm.196604angio-associated migratory protein, re-Mm.147946MYB bind	rin (importin) beta 3 DNA 3010025E17 gene ling protein (P160) 1a I protein S3
Mm. 196581 mitogen activated protein kinase 1 Mm.14663 ATP synt	expression 3 thase, H+ transporting, mi-
	al F0 complex, subunit g c translation initiation factor
like 1 Mm.142740 metallothic	onein 2 protein, large P2
Mm.195894 Mus musculus, clone MGC:11792 Mm.141443 lactate del	protein L9 hydrogenase 1, A chain
Mm.19169 thioredoxin-like (32kD) Mm.140380 ribosomal	ji network protein 2 protein L23
nase (PGK1) betal,4-ga	culus, Similar to xylosylprotein lactosyltransferase, poly- (galactosyltransferase I), clo-
• • • • • • • • • • • • • • • • • • • •	28643 IMAGE:4224150, mR-
·	mouse version of p180 ribo- eptor/ribosome binding prote-
Mm.182931 phosphoribosylaminoimidazole car- 30 Mm.13020 ribosomal boxylase, phosphoribosylaminoribosy- laminoimidazole, succinocarboxamide binding, fa	protein L13a peta (A4) precursor protein- amily A, member 3
- · · · · · · · · · · · · · · · · · · ·	in (importin) alpha 2
body cDNA, RIKEN full-length enriched Mm.11376 ribosomal library, clone:5730409M10:CCAAT/en- Mm.1125 expressed	amma subunit (S. cerevisiae) protein L36 I in non-metastatic cells 2, IM23B) (nucleoside diphos-
quence quence 40 Mm.1120 endometria Mm.181740 interferon-related developmental regu- Mm.108076 phosphofra	al bleeding associated factor uctokinase, platelet
	NA 2010004J23 gene nouse version of 60S riboso- n I 4
Mm.17932 purine-nucleoside phosphorylase 45 Mm.10702 calcyclin b Mm.1777 heat shock protein, 60 kDa Mm.10665 Mus muscu	inding protein ulus, clone IMAGE:3498496,
Mm.175848 (Manual) small Ca-binding protein Cal- Mm.10623 expressed	sequence Al480570 dehydrogenase
TIDE) (EMAP) Mm.10474 RIKEN cDI	rier family 1, member 7 NA 3110005M08 gene
Mm.175661RIKEN cDNA 1110036C17 geneMm.101619ESTMm.1710hydroxymethylbilane synthaseMm.10spermidineMm.17031POU domain, class 5, transcription fac-Mm.4325Kruppel-lik	e synthase se factor 4 (gut) [Swissprot:
tor 1 55 splQ60793	g;splQ9R255;] growth factor 2, binding pro-
Mm.1639 myeloid cell leukemia sequence 1 tein 1 [Swis	ssprot: splO88477;] 2 [Swissprot: splO88322;

_	_
"	۰,

	splQ8R5G0;splQ9CT94;]		Mm.7793	protein phosphatase 1, catalytic subu-
Mm.34407	MAD homolog 7 (Drosophila) [Swiss-			nit, gamma isoform
	prot: splO35253;splQ9CSC7;]		Mm.7723	poly(A) binding protein, nuclear 1
Mm.4451	hairy and enhancer of split 1, (Droso-		Mm.76278	RIKEN cDNA 2610203K23 gene
	phila) [Swissprot: none]	5	Mm.7516	nuclear autoantigenic sperm protein (hi-
Mm.57195	nodal [Swissprot: spIP43021;]		** 7040	stone-binding)
Mm.1249	laminin, gamma 1 [Swissprot: spl		Mm.7312	DNA segment, Chr 17, human D6S56E 2
Mm.27706	P02468;] ash2 (absent, small, or homeotic)-like		Mm.7141	proliferating cell nuclear antigen
141111.27700	(Drosophila) [Swissprot:	10	Mm.6787	splicing factor, arginine/serine-rich 3
Mm.4603	scavenger receptor class B1 [Swiss-			(SRp20)
	prot: splQ61009;splQ9CWJ7;}		Mm.66	ribosomal protein S4, X-linked
Mm.181562	adhesion regulating molecule 1 [Swis-		Mm.6476	RIKEN cDNA 2700084L22 gene
	sprot: splQ8VCl8;splQ922A7;		Mm.64104	RIKEN cDNA 2410016F19 gene
	splQ9JKV1;]	15	Mm.6343	nucleophosmin 1
Mm.43444	MAD2 (mitotic arrest deficient, homo-		Mm.61901	expressed sequence Al429604
M 100075	log)-like 1 (yeast) [Swissprot:		Mm.6065	inosine 5'-phosphate dehydrogenase 2
Mm.103675 Mm.980	sacsin [Swissprot: none] tenascin C [Swissprot: splQ64706;		Mm.5624	DEAD/H (Asp-Glu-Ala-Asp/His) box po-
WITH. 960	splQ9WUU4;]	20	Mm.548	lypeptide 16 cytochrome c oxidase, subunit VIc
Mm.5090	cripto, teratocarcinoma-derived growth		Mm.5305	guanine nucleotide binding protein, beta
	factor (Tdgf1)			2, related sequence 1
Mm.30177	D11Ertd603e, DNA segment, Chr 11,		Mm.525	eukaryotic translation initiation factor 4,
	ERATO Doi 603			gamma 2
Mm.233844	C330012H03Rik, RIKEN cDNA	25	Mm.5114	dishevelled 2, dsh homolog (Drosophi-
	C330012H03			la)
	Normania a constanti di Chamana di Constanti		Mm.4933	mini chromosome maintenance de-
Liste E: Posi	tivmarker neurale Stammzellen (2.);		Mm.4890	ficient 6 (S. cerevisiae) Finkel-Biskis-Reilly murine sarcoma vi-
FS-7ellen ±	PSA-NCAM -/+; Adult brain -	30	WIII1.4690	rus (FBR-MuSV) ubiquitously expres-
LO Lenen +,	ON-ITOMIT - TT, Madic Blain -	••		sed (fox derived)
[0043]			Mm.4846	lamin B1
•			Mm.4756	leptin receptor
Mm.99776	cytosolic aminopeptidase P			
	cytosolic animopephidase r		Mm.46754	expressed sequence Al316867
Mm.9916	RIKEN cDNA 1200008012 gene	35	Mm.46754 Mm.46533	RIKEN cDNA 1110007L15 gene
Mm.99	RIKEN cDNA 1200008012 gene ribonucleotide reductase M2	35	Mm.46533 Mm.4551	RIKEN cDNA 1110007L15 gene villin 2
Mm.99 Mm.9811	RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene	35	Mm.46533	RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1
Mm.99	RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corre-	35	Mm.46533 Mm.4551 Mm.4550	RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide
Mm.99 Mm.9811	RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corresponding to human splQ9Y3l0, similar		Mm.46533 Mm.4551 Mm.4550 Mm.4541	RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2
Mm.99 Mm.9811 Mm.9257	RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corresponding to human splQ9Y3l0, similar to E.coli rtcB, UPF0027-family	<i>35</i>	Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312	RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5
Mm.99 Mm.9811	RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corresponding to human splQ9Y3l0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1		Mm.46533 Mm.4551 Mm.4550 Mm.4541	RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2
Mm.99 Mm.9811 Mm.9257	RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corresponding to human splQ9Y3l0, similar to E.coli rtcB, UPF0027-family		Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149	RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs
Mm.99 Mm.9811 Mm.9257	RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corresponding to human splQ9Y3l0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-re-		Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132	RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759
Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918	RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corresponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-regulated protein, 78kD) high mobility group nucleosomal binding domain 2		Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.43444	RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast)
Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918	RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corresponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-regulated protein, 78kD) high mobility group nucleosomal binding domain 2 heterogeneous nuclear ribonucleopro-	40	Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.43444 Mm.4280	RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene
Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911	RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corresponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-regulated protein, 78kD) high mobility group nucleosomal binding domain 2 heterogeneous nuclear ribonucleoprotein L	40	Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.43444 Mm.4280 Mm.42767	RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17
Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927	RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corresponding to human splQ9Y3I0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-regulated protein, 78kD) high mobility group nucleosomal binding domain 2 heterogeneous nuclear ribonucleoprotein L signal recognition particle 9 kDa	40	Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.43444 Mm.4280 Mm.42767 Mm.4237	RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha
Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579	RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corresponding to human splQ9Y3l0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-regulated protein, 78kD) high mobility group nucleosomal binding domain 2 heterogeneous nuclear ribonucleoprotein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1	40	Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.43444 Mm.4280 Mm.42767	RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha proteasome (prosome, macropain) sub-
Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579 Mm.89136	RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corresponding to human splQ9Y3l0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-regulated protein, 78kD) high mobility group nucleosomal binding domain 2 heterogeneous nuclear ribonucleoprotein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1 H3 histone, family 3A	40 45	Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.43444 Mm.4280 Mm.42767 Mm.4237 Mm.42197	RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha
Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579	RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corresponding to human splQ9Y3l0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-regulated protein, 78kD) high mobility group nucleosomal binding domain 2 heterogeneous nuclear ribonucleoprotein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1	40 45	Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.43444 Mm.4280 Mm.42767 Mm.4237	RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha proteasome (prosome, macropain) subunit, beta type 1
Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579 Mm.89136 Mm.88212	RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corresponding to human splQ9Y3l0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-regulated protein, 78kD) high mobility group nucleosomal binding domain 2 heterogeneous nuclear ribonucleoprotein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1 H3 histone, family 3A tubulin, alpha 6 mammary tumor integration site 6 baculoviral IAP repeat-containing 5	40 45	Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.4244 Mm.4280 Mm.42767 Mm.4237 Mm.42197	RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha proteasome (prosome, macropain) subunit, beta type 1 catalase 1
Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579 Mm.89579 Mm.89136 Mm.88212 Mm.880	RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corresponding to human splQ9Y3l0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-regulated protein, 78kD) high mobility group nucleosomal binding domain 2 heterogeneous nuclear ribonucleoprotein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1 H3 histone, family 3A tubulin, alpha 6 mammary tumor integration site 6	40 45	Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.4280 Mm.42767 Mm.4237 Mm.42197 Mm.4215 Mm.41940 Mm.4189 Mm.41023	RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha proteasome (prosome, macropain) subunit, beta type 1 catalase 1 RIKEN cDNA 6530409L22 gene cyclin A2 RIKEN cDNA 1110021E09 gene
Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579 Mm.89136 Mm.88212 Mm.880 Mm.8552 Mm.8552	RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corresponding to human splQ9Y3l0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-regulated protein, 78kD) high mobility group nucleosomal binding domain 2 heterogeneous nuclear ribonucleoprotein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1 H3 histone, family 3A tubulin, alpha 6 mammary tumor integration site 6 baculoviral IAP repeat-containing 5 KH domain containing, RNA binding, signal transduction associated 1	40 45	Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.4280 Mm.4280 Mm.42767 Mm.4237 Mm.42197 Mm.4215 Mm.41940 Mm.4189	RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha proteasome (prosome, macropain) subunit, beta type 1 catalase 1 RIKEN cDNA 6530409L22 gene cyclin A2 RIKEN cDNA 1110021E09 gene antigen identified by monoclonal antibo-
Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579 Mm.89579 Mm.88552 Mm.8852 Mm.8552 Mm.8552	RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corresponding to human splQ9Y3l0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-regulated protein, 78kD) high mobility group nucleosomal binding domain 2 heterogeneous nuclear ribonucleoprotein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1 H3 histone, family 3A tubulin, alpha 6 mammary tumor integration site 6 baculoviral IAP repeat-containing 5 KH domain containing, RNA binding, signal transduction associated 1 TG interacting factor	40 45	Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.4280 Mm.42767 Mm.42767 Mm.42197 Mm.4215 Mm.4189 Mm.41023 Mm.41023 Mm.4078	RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha proteasome (prosome, macropain) subunit, beta type 1 catalase 1 RIKEN cDNA 6530409L22 gene cyclin A2 RIKEN cDNA 1110021E09 gene antigen identified by monoclonal antibody Ki 67
Mm.99 Mm.9811 Mm.9257 Mm.925 Mm.918 Mm.911 Mm.9043 Mm.89927 Mm.89579 Mm.89136 Mm.88212 Mm.880 Mm.8552 Mm.8552	RIKEN cDNA 1200008012 gene ribonucleotide reductase M2 RIKEN cDNA 2310008M10 gene (Manual) uncharacterized protein corresponding to human splQ9Y3l0, similar to E.coli rtcB, UPF0027-family transcription factor Dp 1 heat shock 70kD protein 5 (glucose-regulated protein, 78kD) high mobility group nucleosomal binding domain 2 heterogeneous nuclear ribonucleoprotein L signal recognition particle 9 kDa mannose-P-dolichol utilization defect 1 H3 histone, family 3A tubulin, alpha 6 mammary tumor integration site 6 baculoviral IAP repeat-containing 5 KH domain containing, RNA binding, signal transduction associated 1	40 45	Mm.46533 Mm.4551 Mm.4550 Mm.4541 Mm.45312 Mm.45149 Mm.45132 Mm.4426 Mm.4280 Mm.42767 Mm.4237 Mm.42197 Mm.4215 Mm.41940 Mm.4189 Mm.41023	RIKEN cDNA 1110007L15 gene villin 2 ATPase, Na+/K+ transporting, beta 1 polypeptide SRY-box containing gene 2 anaphase-promoting complex subunit 5 ESTs expressed sequence AW121759 Cd63 antigen MAD2 (mitotic arrest deficient, homolog)-like 1 (yeast) RIKEN cDNA 2010203J19 gene ribosomal protein S17 topoisomerase (DNA) II alpha proteasome (prosome, macropain) subunit, beta type 1 catalase 1 RIKEN cDNA 6530409L22 gene cyclin A2 RIKEN cDNA 1110021E09 gene antigen identified by monoclonal antibo-

Mm.4024 Mm.3931	cofilin 1, non-muscle Max protein		Mm.29122 Mm.29055	RIKEN cDNA 0610012D09 gene chromobox homolog 1 (Drosophila HP1
Mm.38930	expressed sequence AA407558			beta)
Mm.38912	RIKEN cDNA 2410129H14 gene	-	Mm.29054	RIKEN cDNA 2610529112 gene
Mm.38611	RIKEN cDNA 2210021A15 gene	5	Mm.29005	expressed sequence AU021749
Mm.38528	RIKEN cDNA 2810430M08 gene		Mm.28995	RIKEN cDNA 2010009J12 gene
Mm.38306 Mm.3797	macrophage erythroblast attacher		Mm.28985 Mm.28965	ribosomal protein L27
Mm.37835	nucleosome assembly protein 1-like 1 ribosomal protein L7		Mm.28964	RIKEN cDNA 0710007A14 gene Mus musculus, clone IMAGE:4949762,
Mm.372	ribosomal protein S26	10	Willi.20904	mRNA, partial cds
Mm.36511	mitochondrial ribosomal protein L32		Mm.28961	cleavage and polyadenylation specific
Mm.35844	growth arrest specific 5		141111.20001	factor 5, 25 kD subunit
Mm.35829	erythroid differentiation regulator		Mm.28909	protein tyrosine phosphatase 4a1
Mm.35661	Mus musculus, Similar to hypothetical		Mm.28899	RIKEN cDNA 1110059P08 gene
	protein, clone MGC:29235 IMAGE:	15	Mm.28805	SET translocation
	5043282, mRNA, complete cds		Mm.28805	SET translocation
Mm.35087	expressed sequence AA673488		Mm.28805	SET translocation
Mm.3501	kinesin family member C5A		Mm.28726	EST C77032
Mm.34914	ESTs		Mm.28694	RIKEN cDNA 2410088K19 gene
Mm.3487	ribosomal protein L30	20	Mm.28560	Ly1 antibody reactive clone
Mm.3444	bromodomain-containing 2		Mm.28499	Mus musculus, similar to CG15881 ge-
Mm.34385	expressed sequence Al450344			ne product (H. sapiens), clone MGC:
Mm.34261	expressed sequence AW557761			36308 IMAGE:5040108, mRNA, com-
Mm.3381	ribosomal protein S8			plete cds
Mm.3380	kinesin family member 5B	25	Mm.28299	ESTs, Highly similar to GUAA_HUMAN
Mm.3360	tyrosine 3-monooxygenase/tryptophan			GMP SYNTHASE
	5-monooxygenase activation protein,		Mm.28222	RIKEN cDNA 2610307C23 gene
14 000	zeta polypeptide		Mm.28121	RIKEN cDNA 1110061A19 gene
Mm.326	RIKEN cDNA 1110038L14 gene		Mm.28044	filamin-like protein
Mm.320	DNA polymerase alpha 2, 68 kDa	30	Mm.27972	NS1-associated protein 1
Mm.3199	RIKEN cDNA 1500001N04 gene		Mm.27927	heterogeneous nuclear ribonucleopro-
Mm.31512 Mm.31228	ring finger protein 2 RIKEN cDNA 1810022K09 gene		Mm.27852	tein A1 expressed sequence AW555814
Mm.30806	ribosomal protein L19		Mm.27818	eukaryotic translation elongation factor
Mm.3054	alpha-L-iduronidase	35	141111.27010	2
Mm.3035	RIKEN cDNA 3110006P09 gene		Mm.27796	RIKEN cDNA 5730427N09 gene
Mm.30270	proteasome (prosome, macropain) sub-		Mm.27669	small nuclear ribonucleoprotein E
	unit, alpha type 4		Mm.27660	RIKEN cDNA 5730420G12 gene
Mm.30120	ribosomal protein S27-like		Mm.27624	RIKEN cDNA C530002L11 gene
Mm.30069	RIKEN cDNA 1200003J11 gene	40	Mm.27293	RIKEN cDNA 4833420K19 gene
Mm.30011	ribosomal protein S23		Mm.27269	RIKEN cDNA 2310037I24 gene
Mm.29931	cell division cycle 20 homolog (S. cere-		Mm.27141	Rac GTPase-activating protein 1
	visiae)		Mm.27074	RIKEN cDNA 2610019N13 gene
Mm.29923	SMT3 (supressor of mif two, 3) homolog		Mm.265	ribosomal protein S25
	2 (S. cerevisiae)	45	Mm.2591	RNA binding motif protein 3
Mm.29911	RIKEN cDNA 3200001M24 gene		Mm.25558	RIKEN cDNA 2410018J24 gene
Mm.29896	ribosomal protein L21		Mm.25542	(Manual) strange EST contig in intron of
Mm.2986	expressed sequence AW146116			p137 (GPI-anchored transcytosis prote-
Mm.29829	expressed sequence Al326010			in), maybe alternative C-terminus of
Mm.29666	solute carrier family 25 (mitochondrial	50	NA OE 4	splQ60865
	carnitine/acylcarnitine translocase),		Mm.254	tumor protein, translationally-controlled
Mm 2066	member 20 ,		Mm 25200	1 ESTs Wookly similar to simple report
Mm.2966	isocitrate dehydrogenase 2 (NADP+), mitochondrial		Mm.25299	ESTs, Weakly similar to simple repeat sequence-containing transcript
Mm.29296	RIKEN cDNA 1110003H02 gene	55	Mm.25164	gene trap locus 1-13
Mm.29194	RIKEN cDNA 1700094M07 gene		Mm.25137	RIKEN cDNA 2410004B18 gene
Mm.29133	budding uninhibited by benzimidazoles		Mm.24870	(Manual assignment) UBP7 ubiquitin
	1 homolog, beta (S. cerevisiae)			hydrolase
	<u> </u>			•

Mm.24591	expressed sequence AW546279		Mm.21054	nuclease sensitive element binding pro-
Mm.2424	ribosomal protein L10A			tein 1
Mm.24219	RIKEN cDNA 1810037I17 gene		Mm.20927	transforming growth factor beta 1 indu-
Mm.24042	RIKEN cDNA 1210001E11 gene			ced transcript 4
Mm.23943	vesicle-associated membrane protein,	5	Mm.206399	ESTs
	associated protein A (33 kDa)		Mm.2038	Ras-GTPase-activating protein SH3-
Mm.23758	RIKEN cDNA 1110008P04 gene			domain binding protein
Mm.23695	dihydrofolate reductase		Mm.2025	survival motor neuron
Mm.23692	casein kinase II, alpha 1 related se-	40	Mm.200837	Mus musculus, clone IMAGE:5355658,
Mm 22006	quence 4	10	M 400044	mRNA
Mm.23096	protein phosphatase 2 (formerly 2A), regulatory subunit B", alpha		Mm.196614	eukaryotic translation elongation factor 1 alpha 1
Mm.2287	proteasome (prosome, macropain) sub-		Mm.196608	expressed sequence AA407306
	unit, alpha type 5		Mm.196526	ADP-ribosylation factor 6
Mm.22731	integrin beta 4 binding protein	15	Mm.196515	DNA segment, Chr 1, ERATO Doi 692,
Mm.2265	U1 small nuclear ribonucleoprotein 1C			expressed
Mm.22387	expressed sequence Al314668		Mm.196396	tubulin, alpha 1
Mm.22269	exportin 1, CRM1 homolog (yeast)		Mm.196365	RIKEN cDNA 4833416109 gene
Mm.22214	RIKEN cDNA 2610008F03 gene		Mm.196328	RIKEN cDNA 5830466J11 gene
Mm.220918	heterogeneous nuclear ribonucleopro-	20	Mm.195898	phosphatidylethanolamine binding pro-
	tein D-like			tein
Mm.220342	Mus musculus, clone IMAGE:3669867,		Mm.1951	ribonucleic acid binding protein S1
M 040070	mRNA, partial cds		Mm.1948	t-complex testis expressed 1
Mm.219670	Mus musculus, Similar to eukaryotic	25	Mm.193688	RIKEN cDNA 2700059D21 gene
	translation initiation factor 4 gamma, 1, clone IMAGE:4950789, mRNA, partial	25	Mm.19187 Mm.19101	prothymosin alpha DEAD (aspartate-glutamate-alanine-
	cds		WIIII. 19101	aspartate) box polypeptide 5
Mm.219668	RIKEN cDNA 2610209F03 gene		Mm.19015	serine racemase
Mm.219648	Mus musculus, Similar to nuclear matrix		Mm.18923	mini chromosome maintenance de-
	protein p84, clone MGC:28284 IMAGE:	30		ficient 7 (S. cerevisiae)
	4010605, mRNA, complete cds		Mm.18921	valosin containing protein
Mm.21964	Mus musculus, clone IMAGE:3485208,		Mm. 18856	mitogen-activated protein kinase 6
	mRNA, partial cds		Mm.18705	vacuolar protein sorting 4b (yeast)
Mm.21873	retroviral integration site 1		Mm.18700	RIKEN cDNA 1200009K13 gene
Mm.218657	cerebellar ataxia 3	35	Mm.18637	teratocarcinoma expressed, serine rich
Mm.21841	splicing factor, arginine/serine-rich 2		Mm.18516	H3 histone, family 3B
M 010040	(SC-35)		Mm.1843	heat shock protein, 86 kDa 1
Mm.218240	Mus musculus, clone IMAGE:5342828,		Mm.183102	actin-related protein 3 homolog (yeast)
Mm.2180	mRNA, partial cds heat shock protein, 84 kDa 1	40	Mm.183016 Mm.181880	thymine DNA glycosylase RIKEN cDNA 1110007A14 gene
Mm.21764	small nuclear ribonucleoprotein poly-	40	Mm.181562	adhesion regulating molecule 1
	peptide G		Mm.1815	cytidine 5'-triphosphate synthase
Mm.21714	RIKEN cDNA 2410003A14 gene		Mm.180873	RIKEN cDNA 2510019J09 gene
Mm.21559	non-POU-domain-containing, octamer		Mm.180873	(Manual) probably reverse tag of 60S ri-
	binding protein	45		bosomal protein L18a
Mm.213452	Mus musculus, clone IMAGE:5320271,		Mm.180409	ubiquitin-conjugating enzyme E2H
	mRNA, partial cds		Mm.180271	RIKEN cDNA 5630400D24 gene
Mm.213020	(Manual) 60S ribosomal protein L32		Mm.17989	chaperonin subunit 8 (theta)
	(RPL32)		Mm.1777	heat shock protein, 60 kDa
Mm.21295	expressed sequence AW214031	50	Mm.1775	hematological and neurological expres-
Mm.21289	ribosomal protein S12		M 477454	sed sequence 1
Mm.21281 Mm.21185	ring finger protein 4 adaptor-related protein complex AP-3,		Mm.177451 Mm.17330	RIKEN cDNA 5730544L10 gene ESTs
CO 1 1 2.111W	beta 1 subunit		Mm.17330	tropomyosin 3, gamma
Mm.2115	heterogeneous nuclear ribonucleopro-	55	Mm.17300	tubulin, beta 5
	tein U	-	Mm.16976	TAF9 RNA polymerase II, TATA box bin-
Mm.21094	DNA segment, Chr 9, Wayne State Uni-			ding protein (TBP)-associated factor, 32
	versity 138, expressed			kDa

EP 1 529 838 A1

31

Mm.16775	ribosomal protein S24			cogene 1
Mm.16767	heterogeneous nuclear ribonucleoprotein A2/B1			<b>g</b>
Mm.16711	mini chromosome maintenance de- ficient 2 (S. cerevisiae)	5	Pa	atentansprüche
Mm.16525	polo-like kinase homolog, (Drosophila)		1.	Zellpopulation, dadurch gekennzelchnet, dass
Mm.1639	myeloid cell leukemia sequence 1			mindestens 5% der Zellen neurale Vorläuferzellen
Mm.16323	eukaryotic translation initiation factor 4A2			sind, die wenigstens einen der in Liste A oder Liste B aufgeführten Marker aufweisen.
Mm.16323	eukaryotic translation initiation factor	10		Zalla anni latina ala danakan kanakan
Mm.156892	4A2 heterogeneous nuclear ribonucleopro- tein D		2.	Zellpopulation, dadurch gekennzelchnet, dass mindestens 5% der Zellen, neurale Vorläuferzellen sind, die wenigstens zwei, bevorzugt wenigstens 3
Mm.15571	amyloid beta (A4) precursor protein			der in Liste A oder Liste B aufgeführten Marker
Mm.154915	ribosomal protein S29	15		aufweisen.
Mm. 153457	RIKEN cDNA 2810406C15 gene			
Mm.148973	RIKEN cDNA 3010025E17 gene		3.	Zellpopulation, nach mindestens einem der An-
Mm.142872	heterogeneous nuclear ribonucleoprotein K			sprüche 1 bis 2, <b>dadurch gekennzeichnet, dass</b> die neuralen Vorläuferzellen keinen in <b>Liste C</b> auf-
Mm.14245	ribosomal protein, large P2	20		geführten Marker aufweisen.
Mm.14244	ribosomal protein L9			7.0
Mm.142363 Mm.140804	RIKEN cDNA 2810036L13 gene		4.	· ·
WITH. 140004	Mus musculus, guanine nucleotide bin- ding protein (G protein), gamma 5, clone			che 1 bis 3, dadurch gekennzeichnet, dass min- destens 25 % der Zellen neurale Vorläuferzellen
	MGC:8292 IMAGE:3593324, mRNA.	25		sind.
	complete cds			Sillo.
Mm.140380	ribosomal protein L23		5.	Zellpopulation nach mindestes einem der Ansprü-
Mm.13886	suppressor of initiator codon mutations,			che 1 bis 4, dadurch gekennzeichnet, dass es
	related sequence 1 (S. cerevisiae)			sich um eine murine Zellpopulation handelt und/
Mm.133825	RIKEN cDNA 0610010123 gene	30		oder die neuralen Vorläuferzellen aus Hirngewebe
Mm.13356	RIKEN cDNA 3110079L04 gene			erhältlich ist.
Mm.131705	Mus musculus, Similar to single-stran-		_	
	ded DNA binding protein, clone MGC:		6.	Verfahren zur Isolierung einer Zellpopulation nach
	41439 IMAGE: 1314987, mRNA, com-	35		mindestens einem der Ansprüche 1 bis 5 mit folgen-
Mm.12858	plete cds eukaryotic translation initiation factor	33		den Schritten:
	4A1			a) Entnahme einer Probe aus dem Hirn
Mm.12706	Mus musculus, Similar to CG11246 ge-			ay Endiamine emer i robe due deminim
	ne product, clone MGC:8248 IMAGE:			b) Isolieren der neuralen Vorläuferzellen unter
	3591968, mRNA, complete cds	40		Verwendung der angegebenen Marker
Mm.12604	sirtuin 1 ((silent mating type information			
	regulation 2, homolog) 1 (S. cerevisiae)			oder
Mm.12568	expressed sequence AW541137			
Mm.12508	karyopherin (importin) alpha 2			<ul> <li>a) Differenzierung von embryonalen Stamm-</li> </ul>
Mm.12441	expressed sequence AU014645	45		zellen zu neuralen Vorläuferzellen,
Mm.124	thymopoletin			b) Isolieren der neuralen Vorläuferzellen unter
Mm.12236 Mm.12145	zinc finger protein 207			Verwendung der angegebenen Marker
Mm.116989	retinoblastoma binding protein 4 actin-like			oder
Mm.111	poly(rC) binding protein 2	50		odei
Mm.10706	RIKEN cDNA 2010004J23 gene			a) Trans-Differenzierung von adulten, nicht
Mm.10474	RIKEN cDNA 3110005M08 gene			neuralen Stammzellen zu neuralen Vorläufer-
Mm. 10409	golgi autoantigen, golgin subfamily a, 4			zellen,
Mm.103675	sacsin			
Mm.1013	ligase I, DNA, ATP-dependent	55		b) Isolieren der neuralen Vorläuferzellen unter
Mm. 101274	RIKEN cDNA 2010008E23 gene			Verwendung der angegebenen Marker
Mm.10076	mitochondrial ribosomal protein L13			
Mm.16469	Nmycl, neuroblastoma myc-related on-			oder

10

15

20

35

45

 a) Differenzierung von adulten, neuralen Stammzellen zu neuralen Vorläuferzellen,

33

b) Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker

oder

- a) Differenzierung von immortalisierten Zellen zu neuralen Vorläuferzellen,
- b) Isolieren der neuralen Vorläuferzellen unter Verwendung der angegebenen Marker.
- Verwendung mindestens eines Markers ausgewählt aus der Liste A oder Liste B zu Identifizierung oder Isolierung von neuralen Vorläuferzellen.
- Antikörper gegen einen Marker aus der Liste A, B oder C.
- Diagnostikmittel enthaltend mindestens einen, bevorzugt zwei oder mehr Substanzen zur Erkennung der Marker der Liste A, B oder C.
- Arzneimittel enthaltend die Zellpopulation nach einem der Ansprüche 1 bis 5.
- Zellpopulation, dadurch gekennzeichnet, dass mindestens 5% der Zellen neurale Stammzellen sind, die wenigstens einen der in Liste D oder Liste E aufgeführten Marker aufweisen.
- 12. Zellpopulation, dadurch gekennzeichnet, dass mindestens 5% der Zellen neurale Stammzellen sind, die wenigstens zwei, bevorzugt wenigstens 3 der in Liste D oder Liste E aufgeführten Marker aufweisen.
- 13. Zellpopulation, nach mindestens einem der Ansprüche 11 bis 12, dadurch gekennzeichnet, dass die neuralen Stammzellen keinen in Liste A oder Liste C aufgeführten Marker aufweisen.
- 14. Zellpopulation nach mindestens einem der Ansprüche 11-13, dadurch gekennzeichnet, dass mindestens 25% der Zellen neurale Stammzellen sind.
- 15. Zellpopulation nach mindestes einem der Ansprüche 11 bis 14, dadurch gekennzeichnet, dass es sich um eine murine Zellpopulation handelt und/oder die neuralen Stammzellen aus Hirngewebe erhältlich.
- 16. Verfahren zur Isolierung einer Zellpopulation nach mindestens einem der Ansprüche 11 bis 15 mit folgenden Schritten:
  - a) Entnahme einer Probe aus dem Hirn

b) Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker

oder

- a) Differenzierung von embryonalen Stammzellen zu neuralen Stammzellen,
- b) Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker

oder

- a) Trans-Differenzierung von adulten, nicht neuralen Stammzellen zu neuralen Stammzellen.
- b) Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker

oder

- a) De-Differenzierung von adulten, neuralen Vorläuferzellen zu neuralen Stammzellen,
- b) Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker

oder

- a) Differenzierung von immortalisierten Zellen zu neuralen Stammzellen,
- b) Isolieren der neuralen Stammzellen unter Verwendung der angegebenen Marker.
- Antikörper gegen einen Marker aus der Liste D, E, A oder C.
- 18. Diagnostikmittel enthaltend mindestens einen, bevorzugt zwei oder mehr Substanzen zur Erkennung der Marker der Liste D, E, A oder C.
  - Arzneimittel enthaltend die Zellpopulation nach einem der Ansprüche 11 bis 15.



# EUROPÄISCHER TEILRECHERCHENBERICHT

Nummer der Anmeldung

der nach Regel 45 des Europäischen Patentübereinkommens für das weitere Verfahren als europäischer Recherchenbericht gilt

EP 03 02 5506

	EINSCHLÄGIGI	E DOKUMENTE	<del>,</del>	
Kategorie	Kennzeichnung des Dokun der maßgebliche	nents mit Angabe, soweit erforderlich n Teile	Betrifft Anspruch	KLASSIFIKATION DER ANMELDUNG (Int.Cl.7)
X	ARSENIJEVIC YVAN ET multipotent neural the cortex of the a EXPERIMENTAL NEUROL Bd. 170, Nr. 1, Jul Seiten 48-62, XP002 ISSN: 0014-4886 * Seite 52, linke Srechte Spalte, Absarchte Spalte, Absarchte Spalte, Absarchie Spalte, Absar	AL: "Isolation of precursors residing in adult human brain" OGY, 1 2001 (2001-07), 2275728 Spalte, letzter Absatz - atz 1 * spalte, letzter Absatz * spalte, Absatz 2 -	1-6,10	C12N5/06 G01N33/53  RECHERCHIERTE SACHGEBIETE (Int.Cl.7) C12N G01N
Die Reche n einem s der Techn Vollständi Unvollstär Nicht rech	ELLSTÄNDIGE RECHEI ernhenablellung ist der Auffassung, de kolchen Umrang nicht entspricht bzw. ik für diese Ansprüche nicht, bzw. nur g recherchierte Patentansprüche: indig recherchierte Patentansprüche: erchierte Patentansprüche: die Beschränkung der Recherche: ine Ergänzungsblatt C	iß ein oder mehrere Ansprüche, den Vorschrifte entsprechen, daß sinnvolle Ermittlungen über o teilweise, möglich sind.	en des EPÜ len Stand	
	Recherchenort	Abechlußdatum der Recherche		Prüfer
	MÜNCHEN	7. April 2004	Nie	buhr-Ebel, K
X : von t Y : von t ande A : techi O : nichi	TEGORIE DER GENANNTEN DOKU besonderer Bedeutung allein betræht besonderer Bedeutung in Verbindung ren Veröffentichung demelben Kateg nologischer Hintergrund bechriftliche Offenbarung chenlikentur.	MENTEN T: der Erfindung zug E: alberes Patentidok et nach dem Anmeldl mit einer D: in der Anmeldung one L: aus anderen Grün	runde liegende T ument, das jedoo edatum veröffent angeführtes Dok den angeführtes	heorien oder Grundsätze h erst am oder Iloht worden ist urnent Dokument

EPO FORM 1503 03.82 (P04009)

4



# EUROPÄISCHER TEILRECHERCHENBERICHT

Nummer der Anmeldung EP 03 02 5506

	EINSCHLÄGIGE DOKUMENTE	KLASSIFIKATION DER ANMELDUNG (Int.CI.7)	
Kategorie	Kennzeichnung des Dokuments mit Angabe, soweit erforderlich der maßgeblichen Teile	Betrifft Anspruch	,,
X	UCHIDA N ET AL: "Direct isolation of human central nervous system stem cells" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, NATIONAL ACADEMY OF SCIENCE. WASHINGTON, US, Bd. 97, Nr. 26, 19. Dezember 2000 (2000-12-19), Seiten 14720-14725, XP002223508 ISSN: 0027-8424 * Zusammenfassung * * Seite 14722, rechte Spalte, letzter Absatz - Seite 14724, rechte Spalte, Absatz 1 * * Abbildungen 1,2 *	11-16,19	RECHERCHIERTE
X	KANEKO Y ET AL: "MUSASHI1: AN EVOLUTIONALLY CONSERVED MARKER FOR CNS PROGENITOR CELLS INCLUDING NEURAL STEM CELLS" DEVELOPMENTAL NEUROSCIENCE, S. KARGER, BASEL, CH, Bd. 22, Nr. 1/2, 2000, Seiten 139-153, XP001033925 ISSN: 0378-5866 * Zusammenfassung * * Abbildung 5 *	11-16,19	SACHGEBIETE (Int.Cl.7)
	EP 1 354 943 A (NAT INST OF ADVANCED IND SCIEN) 22. Oktober 2003 (2003-10-22) "Monoclonal antibodies, hybridomas, cell isolation method, isolated cells and immunological diagnostic method" * Spalte 2, Zeile 42 - Spalte 3, Zeile 11 * * Spalte 13, Zeile 41 - Spalte 14, Zeile 53 *	1-19	

EPO FORM 1503 03.82 (POAC12)



# EUROPÄISCHER TEILRECHERCHENBERICHT

Nummer der Anmeldung EP 03 02 5506

	EINSCHLÄGIGE DOKUMENTE	KLASSIFIKATION DER ANMELDUNG (Int.Cl.7)	
ategorie	Kennzeichnung des Dokuments mit Angabe, soweit erforderlich der maßgeblichen Teile	Betrifft Anspruch	
X	GIMONA MARIO ET AL: "Beta-Actin Specific Monoclonal Antibody" CELL MOTILITY AND THE CYTOSKELETON, Bd. 27, Nr. 2, 1994, Seiten 108-116, XP009028901 ISSN: 0886-1544 * das ganze Dokument *	8,9,17, 18	RECHERCHIERTE SACHGEBIETE (Int.Cl.7)
			ì

EPO FORM 1503 03.82 (P04C12)



# UNVOLLSTÄNDIGE RECHERCHE ERGÄNZUNGSBLATT C

Nummer der Anmeldung EP 03 02 5506

Unvollständig recherchierte Ansprüche: 6, 16

Grund für die Beschränkung der Recherche (nicht patentfähige Erfindung(en)):

Artikel 52 (4) EPÜ – Verfahren zur chirurgischen Behandlung des menschlichen oder tierischen Körpers

Weitere Beschränkung der Recherche

Unvollständig recherchierte Ansprüche: 1-5, 7-15, 17-19

Grund für die Beschränkung der Recherche:

In den Listen A-E, auf die sich in den Patentansprüchen bezogen wird, sind insgesamt etwa 1000 putative Positiv- und Negativmarker neuraler Vorläuferzellen und neuraler Stammzellen aufgelistet. Diese putativen Marker sind teilweise bereits bekannte Proteine, wie z.B. beta-Aktin oder Interleukin 1 alpha, teilweise aber auch undefinierte, als "ESTs" benannte sogenannte Marker oder partielle mRNA-Sequenzen. Aufgrund der grossen Anzahl der putativen Marker und deren tw. mangelhaften Identifikation ist es unmöglich, eine vollständige Recherche zu erstellen.

## ANHANG ZUM EUROPÄISCHEN RECHERCHENBERICHT ÜBER DIE EUROPÄISCHE PATENTANMELDUNG NR.

EP 03 02 5506

In diesem Anhang sind die Mitglieder der Patentfamilien der im obengenannten europäischen Recherchenbericht angeführten Patentdokumente angegeben. Die Angaben über die Familienmitglieder entsprechen dem Stand der Datei des Europäischen Patentamts am Diese Angaben dienen nur zur Unterrichtung und erfolgen ohne Gewähr.

07-04-2004

ang	lm Recherchenbe eführtes Patentdo	richt kument	Datum der Veröffentlichung		Mitglied(er) Patentfami	der lie	Datum der Veröffentlichung
EP	1354943	A	22-10-2003	EP JP US	1354943 2004002350 2003186335	A2 A A1	22-10-2003 08-01-2004 02-10-2003
						•	

Für nähere Einzelheiten zu diesem Anhang : siehe Amtsblatt des Europäischen Patentamts, Nr.12/82

EPO FORM PO461